

SEPTMEMBER 15, 2011 REPORT
ROBBO HOLLERAN
LIST OF EXHIBITS, TABLES, AND FIGURES

I. LIST OF EXHIBITS

Exhibit 1. (i) Harvest Prescription Fact Sheet; (ii) Management Plan Summary Form - Form 4; (iii) Use Value Appraisal Forest Management Plan - Form 2; (iv) Map, dated and timed 10/21/09 - 11:27:20 AM; and (v) Ortho-Photo Map, dated and timed 10/21/09 - 11:27:20 AM.

Exhibit 2. "Silvicultural Guide for Northern Hardwood Types in the Northeast (revised)" by Dr. William B. Leak, et al., published by the United States Department of Agriculture, Research Paper NE-603 (the "Guide").

Exhibit 3. Robbo Holleran resume and qualifications.

Exhibit 4. November 30, 2011 Decision by Commissioner Sarah Clark (the "Decision Memo").

Exhibit 5. 2006 Use Value Program Manual (the "Program Manual").

Exhibit 6. April 26, 2010 Adverse Inspection Report (the "Report").

Exhibit 7. *Technical Guide to Wildlife Habitat in New England*, DeGraaf, Yamasaki, Leak and Lester, University of Vermont Press (2006), p. 102 -103.

Exhibit 8. Revised map.

II. LIST OF TABLES

Table 1. Summary table of my inventory findings.

Table 2. Stand #24 Plots for Stand Total and All Plots Taken in Stand.

Table 3. Stand #34 Plots Used for Stand Total.

Table 4. Stand #34 Alleged Cut Contrary.

Table 5. Stand #34 Plots Taken in Un-cut Portion.

Table 6. Stand #34 All Plots Taken in Stand.

Table 7. Stand #43 Alleged Cut contrary and Plots Used for Stand Total.

Table 8. Stand #43 All Plots Taken in Stand.

Table 9. Stand #44 All Information.

III. LIST OF FIGURES

Figure 1. Stand #24 Basal Area Per Plot, August 2011.

Figure 2. Stand #34 Basal Area Per Plot, August/September 2011.

Figure 3. Stand #43 Basal Area Per Plot, August/September 2011.

Figure 4. Stand #44 Basal Area Per Plot, August/September 2011.

Exhibit 1.

(i) Harvest Prescription Fact Sheet; (ii) Management Plan Summary Form - Form 4; (iii) Use Value Appraisal Forest Management Plan - Form 2; (iv) Map, dated and timed 10/21/09 - 11:27:20 AM; and (v) Ortho-Photo Map, dated and timed 10/21/09 - 11:27:20 AM.

Amended 10/27/09

HARVEST PRESCRIPTION FACT SHEET

PROJECT CODE: 7086

CONTRACT NUMBER: LM-03-01-09

FORESTER: JCH

TOWN: Lemington

PHOTO: 220264, 220260

STANDS: 24, 34, 43, 46, 54

ROADS: Sims Hill, Upper Clough Brook

JOB DESIGNATION: Upper Clough Brook North

TOTAL ACRES IN STANDS: +/- 471

TOTAL ACRES IN STANDS TO BE TREATED: +/- 471

PROPOSED CONTRACTOR: (TBA)

HARVESTING EQUIPMENT RESTRICTIONS OR REQUIREMENTS: The harvest can be accomplished by a mechanized crew with feller-buncher and grapple skidders or cut-to-length equipment.

CURRENT STAND CONDITION:

Stand 24 is a well stocked Softwood dominated mixed wood type with a total basal area of 110 ft², of that 50 ft² is acceptable growing stock. The stand is dominated by Balsam fir, Red spruce, White birch, Red maple and Yellow birch. The mean stand diameter is 7.9 inches. The stand is weighted towards the medium saw timber size class. The White birch is in severe decline and the majority of Balsam fir is mature. The understory is well stocked with Red spruce seedling and saplings.

Stand 34 is a well stocked Northern hardwood type with a total basal area of 82 ft², of that 35 ft² is acceptable growing stock. The stand is dominated by Sugar maple, Yellow birch, Beech, Balsam fir, Red Spruce. The mean stand diameter is 8.4 inches. The stand is weighted towards the medium saw timber size class. The current stand has a high level of residual stand damage and a fair amount of crown die-back. The understory varies greatly in stocking of acceptable regeneration, with small pockets of Sugar maple seedling and saplings in a patchy distribution around the stand.

Stand 43 is a well stocked Hardwood dominated mixed wood type with a total basal area of 88 ft², of that 38 ft² is acceptable growing stock. The stand is dominated by Yellow birch, Balsam fir, White birch, Red spruce, Red maple. The mean stand diameter is 8.2 inches. The stand is weighted towards the medium saw timber size class. The White birch in the stand is in severe decline and the majority of Balsam fir is mature. The majority of the stand has good stocking in Red spruce seedlings and saplings in the understory.

Stand 44 is a well stocked Northern hardwood type with a basal area of 97 ft², of that 42 ft² is acceptable growing stock. The stand is dominated by Sugar maple, Yellow birch, Beech, Balsam fir, Red spruce. The mean stand diameter is 7.6 inches. The stand is weighed towards the small saw timber size class. The stand has a fair amount of acceptable stocking in the of small saw timber size class.

Stand 46 is a poorly stocked Northern Hardwood type with a basal area of 40 ft², of that 8 ft² is acceptable growing stock. The stand is dominated by Sugar maple, Yellow birch, Beech. The mean stand diameter is 6.8 inches. The stand is weighted towards the large pole timber size class. The stand is well stocked with Sugar maple and Yellow birch seedlings and saplings mixed in with beech and striped maple.

Stand 54 is a moderately stocked Northern hardwood stand with a basal area of 72 ft², of that 28 ft² is acceptable growing stock. The stand is dominated by Sugar maple, Yellow birch, Beech, Red spruce, Balsam fir. The mean stand diameter is 8.2 inches. The stand is weighted towards the medium saw timber size class. The understory is dominated by beech and striped maple.

REGENERATION: Does not include non-commercial species or Beech

Stand	Seedlings per Acre	1"-3" Stems per Acre	Total Stems per Acre	Dominate Acceptable Species
LM-03-24	364	217	581	Red spruce
LM-03-34	198	92	290	Sugar maple
LM-03-43	290	120	410	Red spruce
LM-03-44	128	66	194	Sugar maple
LM-03-46	310	88	398	Sugar maple, Yellow birch
LM-03-54	120	33	153	Sugar maple, Yellow birch

ELEVATION & TERRAIN: Elevation ranges from 1700-2500 ft. with poorly drained soils. The ground is gradual in slope, but increases on the far western side of the harvest unit.

SURROUNDING LANDSCAPE: The unit is located at the end of Sims Hill Road and abuts the Lemington Town forest. The proposed activities are consistent with accepted forest management activities in the region and pose no adverse affect to adjacent biological communities. The harvest unit is not visible to the public.

DESIRED GOAL OF HARVEST:

- Capture value in declining timber
- Retain good quality trees as seed source and retained value for the long term
- Improve stand quality and provide openings for natural regeneration to occur where it is deficient
- Protect and release desirable advanced regeneration
- Protect riparian zones and wetland habitat

RECOMMENDED TREATMENT PRESCRIPTION:

Stand 24 will receive a Two Staged Shelterwood (2SS) (even age UVA code 3) and an Overstory Removal (OSR) (even age UVA code 4). The Shelterwood will target the at-risk and mature stems in areas of the stand that have higher stocking in acceptable growing stock. The target residual basal area is 60 ft². The harvest will release quality growing stock and provide gaps to promote regeneration. The final cut will take place once acceptable levels of regeneration are established. Approximately 30-40% of the stand will receive this treatment. The remaining 60-70% of the stand will receive an Overstory Removal releasing the advanced Red spruce regeneration underneath an unacceptable and mature overstory. Care will be taken during layout to avoid the destruction or damage of regeneration.

Stand 34 will receive a Two Stage Shelterwood (2SS) (even age UVA code 3). The stand lacks an acceptable amount of regeneration and the majority of the overstory is unacceptable growing stock. A low density shelterwood with a residual basal area of 30-40 ft² will be utilized to discourage the establishment of beech in the understory. The shelterwood will be irregular in distribution and will target Sugar maple and Yellow birch with large crowns to provide shade and seed distribution. The portions of the stand will also receive 1-2 acre patches where quality and stocking are not insufficient for a Shelterwood. The patches will not affect the overall stand residual basal area of 30-40 ft².

Stand 43 will receive a Two Staged Shelterwood (2SS) (even age UVA code 3) and Overstory Removal (OSR) (even age UVA code 4). 30-40% of the stand will receive an Overstory Removal where the overstory is in severe decline and the understory is well stocked with seedling and sapling sized Red spruce. The remaining portion of the stand will receive a Shelterwood. The harvest will target the at-risk and mature stems. The target residual basal area is 60 ft². The harvest will release quality growing stock and provide gaps to promote regeneration.

Stand 44 will receive an Intermediate Thinning (ITH) (even age UVA code 2). The stand is well stocked with small saw timber, however many of the medium and large saw timber stems are in decline. The thinning will target the at-risk and mature stems and leave a target residual basal area of 60 ft². This release the small saw timber size class and open up gaps for regeneration.

Stand 46 will receive an Overstory Removal (OSR) (even age UVA code 4). The overstory is in decline and the current stand conditions warrant removal. The understory has good stocking in Sugar maple and Yellow birch seedlings and saplings mixed within Beech and Non-commercial species. Care will be taken during layout to reduce the disturbance of the advanced regeneration. In small areas that do not have sufficient levels of acceptable regeneration a seed tree will be left to ensure regeneration. Yellow birch, Red Spruce or clumps of Sugar maple will be targeted as seed trees.

Stand 54 will receive a Seed Tree Patch cut (STC) (even age UVA code 5). The stand has many pockets of acceptable regeneration established in openings from a previous entry, however small pockets of mature and at risk stems remain and do not have sufficient levels of regeneration to warrant an Overstory Removal. The pockets of remaining overstory will be removed and individual Yellow birch or clumps of Red spruce or Sugar maple will be left to provide seed. Seed Tree Patches will not exceed 25 acres in size across the stand.

Pre-harvest layout will include:

- Identify and protect streams and significant wetlands using blue flagging for the centerline of non-eased stream buffers and the side of the stream for eased buffers and wetlands.
- Identify any potential problems and address them with the contractor.
- Identify former skid trails that can be used in current operations and flag new trails where needed using orange flagging.
- Flag stream crossings will have two pink flags and will designate stream crossing method with written instructions on flagging at crossing location and shown on LV's stream crossing worksheet.
- Require Feller-Buncher to lay all felled wood in skid trails to limit damage to the advanced regeneration and residual stand.
- Minimize skid trails to limit site disturbance.
- Establish bounds of harvest in flagged in pink or pink with "Timber Harvest Boundary" printed in black letters. Where the unit adjoins the property line the unit will be marked in blue flagging.

DESIGNATION OF TIMBER FOR HARVEST:

All stands within this unit (30) will be marked in blue or orange paint and three slashes with a C to indicate that all trees within the paint are to be cut. Marked timber will utilize blue or orange paint. Slashes or spots on at least two sides of the tree and a stump spot indicate removal. Skid trails will be designated with orange flagging. Trees marked with a "W", "X", "LX" or "SB" should not be harvested since their designations refer to wildlife trees (w), no cut trees (x), no cut-line trees (LX) and stream buffer (SB).

STREAM / LEGACY BUFFER PRESCRIPTION:

Larger order "blue line" streams and will have a 50' buffer where no operation will take place and crossings will be very limited. The 50' buffer will also apply to wetlands and other riparian features when appropriate. Blue flagging will be used to identify the buffer on the large order eased streams, and lower order non-eased streams will have blue flagging to designate the estimated stream centerline.

TRUCK ROAD CONDITIONS: No road upgrades are needed

SKID TRAILS:

Trails will be established by LV staff and designated in orange flagging. Any unacceptable trails will be identified as unsuitable and new trails will be put in place where needed.

LANDINGS:

Existing landings will be utilized for the harvest.

REGULATORY CONSIDERATIONS:

//A

PERMIT REQUIREMENTS:

Heavy Cut (Act 15) notification required.

HABITAT RETENTION DESIGNATION

Retention will be utilized to retain native tree and vegetation species for the reoccupation of an implemented clear-cut or overstory removal. The retention will occur in the form of corridor retention along blue line and non blue line streams as well as through standing dead and coarse woody debris recruitment. These stands also feature sensitive hydrological features that will be avoided thus contributing to retention. For information on implementation techniques see pages 51-53 of the Forest Management Plan, Nov. 2007.

THREATENED & ENDANGERED SPECIES CONSIDERATIONS:

N/A

SIGNAGE / NOTIFICATION REQUIREMENTS:

Intent to cut notification to FP&R and VLT due to the CE & UVA

CONSERVATION EASMENT CONDITIONS:

As addressed in page 2 of the Champion Working Forest Conservation Easement it is required that two 16" or greater logs per acre of standing dead or downed trees be established or retained. This includes the recruitment of coarse woody debris by selecting trees that have the possibility of fulfilling the requirement in the future. For more detail see page 54 of the Forest Management Plan, Nov 2007.

CLOSE OUT REQUIREMENTS:

Refer to contract for specs.

Parcel ID For Data Entry (by state)# _____ "FP&R COUNTY I" "ESTER USE ONLY"
 Year of Plan _____ Year of Entry _____
 Year of Last Inspection _____

- 1) Landowner Name (last name, first name) Plum Creek
 2) Landowner Address (Street, PO Box) 46 Mountain Avenue, P.O. Box 400
 (Town) Fairfield (State) ME (Zip Code) 04937
 3) Town That Parcel Is Located In Lemington 4) Total Forestry Acres in Parcel 86,262 (Grand list acreage, minus agricultural or non-productive land and exclusions)
 5) Plan Preparer (last name, first name) Horton, Jonathan 6) Previous Owner (last name, first name) _____
 7) Signature [Signature] 8) Date signed 11/3/09

9) Stand information: (this information is for data entry only and does not override what is in actual plan)

Stand #	Acres	Even-aged ⁽¹⁾ Uneven-aged ⁽²⁾ (existing)	Predominant Site Class (1, 2, 3 or 4)	Timber Type	Quadratic M.S.D.	Total BA	AGS BA	Mgmt. Activities	Scheduled Date (± 3 yrs.)
LM-01-24	68.8	1	1-2	11	7.9	110	50	3/4	2011
LM-01-34	137	1	1-2	06	8.4	82	35	3	2011
LM-01-43	115	1	1-2	11	8.2	88	38	3/4	2011
LM-01-44	37	1	1-2	06	7.6	97	42	2	2011
LM-01-46	16	1	1-2	06	6.8	40	8	4	2011
LM-01-54	97	1	1-2	06	8.2	72	28	5	2011

¹ Update of an existing plan that includes all new stand descriptive data. Generally done on a 10-year cycle.
² Change to an existing plan, generally due to purchase or sale of a portion of the property, or a change in prescription. Does not change the 10-year cycle of the existing plan. If this form is filed with an amendment, indicate the amended information in the appropriate stand, and write an explanation in section 13. Amendments must be signed by the landowner(s).

- 10) No activity - (identify stand # and reasons) _____
- 11) Management Activities - other (identify stand #) _____ 54 will receive a Seed Tree Cut
- 12) Timber Types - other (identify stand #) _____
- 13) Amended prescriptions - (identify stand #) _____

DEFINITIONS	FOREST TYPES	CODE #
ACRES-round off to nearest acre	aspen and/or white birch	01
EVEN-AGED-stands with two or less size classes	white pine, red oak	02
UNEVEN-AGED-stands with three or more size classes	white pine	03
SITE CLASS-predominant site class as defined by UV A standards	hemlock	04
QUADRATIC MEAN STAND DIAMETER - to nearest inch	sugar maple	05
TOTAL BA-basal area to nearest ft ² /acre	beech, birch, sugar maple	06
AGS BA (acceptable growing stock basal area) to nearest ft ² /acre	beech, red maple	07
	spruce	08
	spruce/fir	09
	pioneer species	10
	mixed wood (25%-65% softwood)	11
	other (identify other in section 12)	12

MANAGEMENT ACTIVITY CODES (if one of the following choices reasonably describes the planned management activity, use it. If not, use #13 other and describe the management activity in Section 11. Note these descriptions are for choosing codes only; they are not the silvicultural standards).

- 1. Non-commercial forest stand improvement - A treatment made to improve the composition, structure, quality, or growth of young stands.

EVEN-AGED MANAGEMENT (Two or less distinct size classes)

- 2. Intermediate thinning - Reduce stocking to B level, or remove up to 33% of basal area in densely overstocked stands. AGS basal area shall not be reduced below recommendations of appropriate guide or 60 sq. ft./ac. when no guide is available.
- 3. Shelterwood cut - Stand basal area reduced to between 30 and 70 sq. ft. for hardwoods, 80 to 120 sq. ft. for softwoods.
- 4. Overstory removal cut - Removing overstory after a new featured stand has become established. This applies to two-aged as well as recently regenerated stands. After harvest, residual stand should have minimum of sixty (60) square feet of acceptable growing stock or 350 stems/acre for stands less than six (6) inches MSD.
- 5. Clearcut - Stand basal area reduced below thirty (30) square feet.
- 6. Progressive clearcutting - Removing a portion of a stand in strips or patches, to be followed by similar treatments at predetermined dates until entire area is clearcut. If you choose this option, indicate the total time period from first to last cut (e.g., 30 years) in section 13.

UNEVEN-AGED MANAGEMENT (Three or more distinct size classes) (NOTE: ratio of AGS to total basal area should be maintained or increased through any management activities.)

- 7. Individual Tree Selection - Harvest or improvement cutting that maintains a fairly uniform and continuous crown cover appropriate for regenerating tolerant species.
- 8. Group Selection - Harvest of groups of trees roughly 1/20 to 2 acres in size. Appropriate for regeneration of species of all tolerance groups.

MISCELLANEOUS CHOICES

- 9. Salvage cut - Removing a portion of a stand because of damage or disease. Resulting residual stand may not fit standards in guides.
- 10. Sugarbush thinning - Cut designed with the sole objective of establishing or improving a sugarbush. Cut should follow sugarbush guides.
- 11. Species conversion - Cut done usually in a mixed stand to favor certain species. Treatment should not favor species that are "off site".
- 12. No activity - Indicate reason such as "stand understocked", in section 10.
- 13. Other - In cases where none of the above choices comes reasonably close to describing the management activity, put your own description in section 11.
- 14. Crop Tree Release - Release of timber crop trees on at least 2 sides. For additional information refer to "Crop Tree Management in Eastern Hardwoods". (NA-TP-19-93).

VT Dept. of Forests, Parks & Recreation – USE VALUE APPRAISAL FOREST MANAGEMENT PLAN – Form 2, Page 1

Applicant Name: Plum Creek

Address: 49 Mountain Ave
PO Box 400
Fairfield, Maine 04937

Applicant Phone: 802-334-8402

Applicant e-mail: jhorton@landvest.com

Town: Lemington

Grand List Description: : 86,262

Stand No. LM-03-24	Acres 68.8	Site Class I-II	Access Distance < 1
------------------------------	----------------------	---------------------------	-------------------------------

Stand History: Harvested by Champion Paper Co. approximately 20-25 years ago.

Stand Description
EVEN-AGE X
Or
UNEVEN-AGE _____
Forest Type: SH3B
Stocking Level: B-line
Quadratic MSD: 7.9
Total BA 110 AGS BA 50

Long Range Objective

EVEN-AGE Management
Present Stand Age: 65-70
Rotation Age: 80

Scheduled Treatment
(Type and Year): Shelterwood(3)
Overstory Removal (4) 2011

UNEVEN-AGE MANAGEMENT
(Complete diameter distribution table and/or Q information)
BA 6-10" AGS UGS UGS Cull
BA 12-14" AGS UGS UGS Cull
BA 16"+ AGS UGS UGS Cull
Q Factor: Present Future

UNEVEN-AGE Management
Species Objectives:
Cutting Cycle:
Residual BA:
Diameter Objectives:

Scheduled Treatment
(Type and Year):

Forest Health Conditions: The stand has high residual stand damage. Beech bark *Nectria* complex.

Description of Stand Conditions Stand 24 is a well stocked Softwood dominated mixed wood type with at total basal area of 110 ft², of that 50 ft² is acceptable growing stock. The stand is dominated by Balsam fir, Red spruce, White birch, Red maple and Yellow birch. The mean stand diameter is 7.9 inches. The stand is weighted towards the medium saw timber size class. The White birch is in severe decline and the majority of Balsam fir is mature. The understorey is well stocked with Red spruce seedling and saplings.

Stand Cruise Data: Prism Factor/Plot size 10 No. of Points/Plots Sampled 6 Sampling Date(s): 10/15/09

VT Dept. of Forests, Parks & Recreation – USE VALUE APPRAISAL FOREST MANAGEMENT PLAN – Form 2, Page 1

Applicant Name: Plum Creek

Address: 49 Mountain Ave
PO Box 400
Fairfield, Maine 04937

Applicant Phone: 802-334-8402
Applicant e-mail: jhorton@landvest.com

Town: Lemington

Grand List Description: 86,262

Stand No. LM-03-34	Acres 137	Site Class I-II	Access Distance < 1
------------------------------	---------------------	---------------------------	-------------------------------

Stand History: Harvested by Champion Paper Co. approximately 20-25 years ago.

Stand Description
EVEN-AGE X
Or
UNEVEN-AGE _____
Forest Type: H3B
Stocking Level: B-line
Quadratic MSD: 8.4
Total BA 82 AGS BA 35

Long Range Objective
EVEN-AGE Management
Present Stand Age: 70-75
Rotation Age: 80
Scheduled Treatment
(Type and Year): Shelterwood (3)
2011

Forest Health Conditions: The stand has high residual stand damage. Beech bark *Nectria* complex.

UNEVEN-AGE MANAGEMENT
(Complete diameter distribution table and/or Q information)
BA 6-10" AGS ___ UGS ___ Cull ___
BA 12-14" AGS ___ UGS ___ Cull ___
BA 16"+ AGS ___ UGS ___ Cull ___
Q Factor: Present ___ Future ___

UNEVEN-AGE Management
Species Objectives:
Cutting Cycle:
Residual BA:
Diameter Objectives:
Scheduled Treatment
(Type and Year):

Description of Stand Conditions Stand 34 is a well stocked Northern hardwood type with a total basal area of 82 ft², of that 35 ft² is acceptable growing stock. The stand is dominated by Sugar maple, Yellow birch, Beech, Balsam fir, Red Spruce. The mean stand diameter is 8.4 inches. The stand is weighted towards the medium saw timber size class. The current stand has a high level of residual stand damage and a fair about of crown die-back. The understory varies greatly in stocking of acceptable regeneration, with small pockets of Sugar maple seedling and saplings in a patchy distribution around the stand.

Stand Cruise Data: Prism Factor/Plot size 10 No. of Points/Plots Sampled 8 Sampling Date(s): 10/13/09

VT Dept. of Forests, Parks & Recreation – USE VALUE APPRAISAL FOREST MANAGEMENT PLAN – Form 2, Page 1

Applicant Name: Plum Creek

Address: 49 Mountain Ave

Applicant Phone: 802-334-8402

PO Box 400

Applicant e-mail: jhorton@landvest.com

Fairfield, Maine 04937

Town: Lemington

Grand List Description: : 86,262

Stand No. LM-03-43	Acres 115	Site Class I-II	Access Distance < 1
------------------------------	---------------------	---------------------------	-------------------------------

Stand History: Harvested by Champion Paper Co. approximately 20-25 years ago.

Stand Description

EVEN-AGE X

Or

UNEVEN-AGE _____

Forest Type: H3B

Stocking Level: B-line

Quadratic MSD: 8.2

Total BA 88 **AGS BA** 38

Long Range Objective

EVEN-AGE Management
Present Stand Age: 70-75
Rotation Age: 80

Scheduled Treatment
(Type and Year): Shelterwood (3), OSR (4) 2011

UNEVEN-AGE MANAGEMENT
(Complete diameter distribution table and/or Q information)

BA 6-10" AGS UGS Cull

BA 12-14" AGS UGS Cull

BA 16"+ AGS UGS Cull

Q Factor: Present Future

UNEVEN-AGE Management
Species Objectives:
Cutting Cycle:
Residual BA:
Diameter Objectives:

Scheduled Treatment
(Type and Year):

Description of Stand Conditions Stand 43 is a well stocked Hardwood dominated mixed wood type with a total basal area of 88 ft², of that 38 ft² is acceptable growing stock. The stand is dominated by Yellow birch, Balsam fir, White birch, Red spruce, Red maple. The mean stand diameter is 8.2 inches. The stand is weighted towards the medium saw timber size class. The White birch in the stand is in severe decline and the majority of Balsam fir is mature. The majority of the stand has good stocking in Red spruce seedlings and saplings in the understory.

Stand Cruise Data: Prism Factor/Plot size 10 No. of Points/Plots Sampled 8 Sampling Date(s): 10/13/09

VT Dept. of Forests, Parks & Recreation – USE VALUE APPRAISAL FOREST MANAGEMENT PLAN – Form 2, Page 1
 Applicant Name: Plum Creek
 Address: 49 Mountain Ave
 PO Box 400
 Fairfield, Maine 04937
 Applicant Phone: 802-334-8402
 Applicant e-mail: jhorton@landvest.com

Town: Lemington

Grand List Description: : 86,262

Stand No. LM-03-44	Acres 37	Site Class I-II	Access Distance < 1
-----------------------	-------------	--------------------	------------------------

Stand Description
 EVEN-AGE X
 Or
 UNEVEN-AGE _____

Long Range Objective

EVEN-AGE Management
 Present Stand Age: 70-75
 Rotation Age: 80

Forest Type: H3A
 Stocking Level: B-line

Scheduled Treatment

(Type and Year): Shelterwood (3)
 2011

Quadratic MSD: 7.6

Total BA 97 AGS BA 42

Forest Health Conditions: The stand has high residual stand damage. Beech bark *Nectria* complex.

UNEVEN-AGE MANAGEMENT
 (Complete diameter distribution table and/or Q information)

BA 6-10" AGS UGS Cull
 BA 12-14" AGS UGS Cull
 BA 16"+ AGS UGS Cull

UNEVEN-AGE Management
 Species Objectives:
 Cutting Cycle:
 Residual BA:
 Diameter Objectives:

Q Factor: Present Future

Scheduled Treatment
 (Type and Year):

Description of Stand 44 is a well stocked Northern hardwood type with a basal area of 97 ft², of that 42 ft² is acceptable growing stock. The stand is dominated by Sugar maple, Yellow birch, Beech, Balsam fir, Red spruce. The mean stand diameter is 7.6 inches. The stand is weighed towards the small saw timber size class. The stand has a fair amount acceptable stocking in the of small saw timber size class.

Stand Cruise Data: Prism Factor/Plot size 10 No. of Points/Plots Sampled 5 Sampling Date(s): 10/13/09

VT Dept. of Forests, Parks & Recreation – USE VALUE APPRAISAL FOREST MANAGEMENT PLAN – Form 2, Page 1

Applicant Name: Plum Creek

Address: 49 Mountain Ave
PO Box 400
Fairfield, Maine 04937

Applicant Phone: 802-334-8402
Applicant e-mail: jhorton@landvest.com

Town: Lemington

Grand List Description: : 86,262

Stand No. LM-03-46	Acres 16	Site Class I-II	Access Distance < 1
------------------------------	--------------------	---------------------------	-------------------------------

Stand History: Harvested by Champion Paper Co. approximately 20-25 years ago.

Long Range Objective

EVEN-AGE Management
Present Stand Age: 60-70
Rotation Age: 80

Scheduled Treatment
(Type and Year): OSR (4) 2011

UNEVEN-AGE Management
Species Objectives:
Cutting Cycle:
Residual BA:
Diameter Objectives:

Scheduled Treatment
(Type and Year):

Stand Description

EVEN-AGE **UNEVEN-AGE**

Or

Forest Type: H3C

Stocking Level: C-line

Quadratic MSD: 6.8

Total BA 40 **AGS BA** 8

UNEVEN-AGE MANAGEMENT

(Complete diameter distribution table and/or Q information)

BA 6-10" **AGS** **UGS** **Cull**

BA 12-14" **AGS** **UGS** **Cull**

BA 16"+ **AGS** **UGS** **Cull**

Q Factor: Present Future

Description of Stand Conditions Stand 46 is a poorly stocked Northern Hardwood type with a basal area of 40 ft², of that 8 ft² is acceptable growing stock. The stand is dominated by Sugar maple, Yellow birch, Beech. The mean stand diameter is 6.8 inches. The stand is weighted towards the large pole timber size class. The stand is well stocked with Sugar maple and Yellow birch seedlings and saplings mixed in with beech and striped maple.

Stand Cruise Data: Prism Factor/Plot size 10 No. of Points/Plots Sampled 3 Sampling Date(s): 10/13/09

Applicant Name: Plum Creek
Address: 49 Mountain Ave
 PO Box 400
 Fairfield, Maine 04937
Applicant Phone: 802-334-8402
Applicant e-mail: jhorton@landvest.com

Town: Lemington
Grand List Description: : 86,262

Stand No. LM-03-54	Acres 97	Site Class I-II	Access Distance < 1
------------------------------	--------------------	---------------------------	-------------------------------

Stand History: Harvested by Champion Paper Co. approximately 20-25 years ago.

Stand Description
 EVEN-AGE
 Or
 UNEVEN-AGE
Forest Type: H3B
Stocking Level: B-line
Quadratic MSD: 8.2
Total BA 72 AGS BA 28

Long Range Objective
 EVEN-AGE Management
 Present Stand Age: 65-70
 Rotation Age: 80
Scheduled Treatment
 (Type and Year): Seed Tree (5) 2011

UNEVEN-AGE MANAGEMENT
 Species Objectives:
 Cutting Cycle:
 Residual BA:
 Diameter Objectives:
Scheduled Treatment
 (Type and Year):

Forest Health Conditions: The stand has high residual stand damage. Beech bark *Nectria* complex.

Q Factor: Present ___ Future ___
 BA 6-10" ___ AGS ___ UGS ___ Cull ___
 BA 12-14" ___ AGS ___ UGS ___ Cull ___
 BA 16"+ ___ AGS ___ UGS ___ Cull ___


Description of Stand 54 is a moderately stocked Northern hardwood stand with a basal area of 72 ft², of that 28 ft² is acceptable growing stock. The stand is dominated by Sugar maple, Yellow birch, Beech, Red spruce, Balsam fir. The mean stand diameter is 8.2 inches. The stand is weighted towards the medium saw timber size class. The understory is dominated by beech and striped maple.

Stand Cruise Data: Prism Factor/Plot size ___ 10 ___ No. of Points/Plots Sampled ___ 6 ___ Sampling Date(s): 10/15/09

**MANAGEMENT SCHEDULE
USE VALUE APPRAISAL FOREST MANAGEMENT PLAN**

STAND NO.	YEAR	MANAGEMENT PRACTICES TO BE ACCOMPLISHED DURING NEXT 10 YEAR PLAN:	Silvicultural Guide or Tech Ref. Prescription #/Letter, if appropriate
LM-03-24	2011	Stand 24 will receive a Two Staged Shelterwood (2SS) (even age UVA code 3) and an Overstory Removal (OSR) (even age UVA code 4). The Shelterwood will target the at-risk and mature stems in areas of the stand that have higher stocking in acceptable growing stock. The target residual basal area is 60 ft ² . The harvest will release quality growing stock and provide gaps to promote regeneration. The final cut will take place once acceptable levels of regeneration are established. Approximately 30-40% of the stand will receive this treatment. The remaining 60-70% of the stand will receive an Overstory Removal releasing the advanced Red spruce regeneration underneath an unacceptable and mature overstory. Care will be taken during layout to avoid the destruction or damage of regeneration.	Guide:
34	2011	Stand 34 will receive a Two Stage Shelterwood (2SS) (even age UVA code 3). The stand lacks an acceptable amount of regeneration and the majority of the overstory is unacceptable growing stock. A low density shelterwood with a residual basal area of 30-40 ft ² will be utilized to discourage the establishment of beech in the understory. The shelterwood will be irregular in distribution and will target Sugar maple and Yellow birch with large crowns to provide shade and seed distribution. The portions of the stand will also receive 1-2 acre patches where quality and stocking are not sufficient for a Shelterwood. The patches will not affect the overall stand residual basal area of 30-40 ft ² .	Guide:
43	2011	Stand 43 will receive a Two Staged Shelterwood (2SS) (even age UVA code 3) and Overstory Removal (OSR) (even age UVA code 4). 30-40% of the stand will receive an Overstory Removal where the overstory is in severe decline and the understory is well stocked with seedling and sapling sized Red spruce. The remaining portion of the stand will receive a Shelterwood. The harvest will target the at-risk and mature stems. The target residual basal area is 60 ft ² . The harvest will release quality growing stock and provide gaps to promote regeneration.	

STAND NO.	YEAR	MANAGEMENT PRACTICES TO BE ACCOMPLISHED DURING NEXT 10 YEAR PLAN:	Silvicultural Guide or Tech Ref. Prescription #/Letter, if appropriate
44	2011	Stand 44 will receive an Intermediate Thinning (ITH) (even age UVA code 2). The stand is well stocked with small saw timber, however many of the medium and large saw timber stems are in decline. The thinning will target the at-risk and mature stems and leave a target residual basal area of 60 ft ² . This release the small saw timber size class and open up gaps for regeneration.	Guide:
46	2011	Stand 46 will receive and Overstory Removal (OSR) (even age UVA code 4). The overstory is in decline and the current stand conditions warrant removal. The understory has good stocking in Sugar maple and Yellow birch seedlings and saplings mixed within Beech and Non-commercial species. Care will be taken during layout to reduce the disturbance of the advanced regeneration. In small areas that do not have sufficient levels of acceptable regeneration a seed tree will be left to ensure regeneration. Yellow birch, Red Spruce or clumps of Sugar maple will be targeted as seed trees.	Guide:
54	2011	Stand 54 will receive a Seed Tree Patch cut (STC) (even age UVA code 5). The stand has many pockets of acceptable regeneration established in openings from a previous entry, however small pockets of mature and at risk stems remain and do not have sufficient levels of regeneration to warrant an Overstory Removal. The pockets of remaining overstory will be removed and individual Yellow birch or clumps of Red spruce or Sugar maple will be left to provide seed. Seed Tree Patches will not exceed 25 acres in size across the stand.	

LANDOWNER'S SIGNATURE:  DATE: 11/13/09
 PREPARED BY: Jonathan Horton DATE: 10/14/09
 CERTIFIED BY: _____ DATE: _____


568000 568500 569000 569500 570000 570500

VT STATE PLANE
NAD 83 METERS



**LAND OF PLUM CREEK MAINE
TIMBERLANDS, LLC.
NORTHERN KINGDOM UNIT-VT
HARVEST MAP
CLOUGH BROOK NORTH
LM-03-01-09**

SCALE 1:15,840 1"=1,320'
0 660 1,320 2,640 Feet



LandVest
Town: LEMINGTON
County: ESSEX
Compartment: LM-03
Forester: JCH
Job Number: LM-03-01-09
Tractor: TBD

STAND_NUMB	STRATA	OS TYPE	SILV PRES	ACRES
LM-03-24	SH34B/SH2B	SH3B	ITH	68.8
LM-03-34.0	H3B	H3B	2SS	137.0
LM-03-43.0	HS3B	HS3B	OSR/ITH	115.1
LM-03-44	H3B	H3A	ITH	37.0
LM-03-46	H3C/H1B	H3C	OSR	15.9
LM-03-54	H3B	H3B	STC	97.4
TOTAL (Acres derived from GIS)				471.1

- PROPOSED HARVEST
- ACTIVE HARVEST
- COMPLETE HARVEST
- APPROVED HARVEST
- POSTPONED HARVEST
- FLOWAGE/BOG
- GRAVEL PIT
- OPEN/AGRICULTURAL LANDING
- RIPARIAN ZONE
- SPECIAL TREATMENT AREA
- WATER
- COUNTY LINE
- TOWN LINE
- STREAM
- STAND LINE
- COMPARTMENT LINE
- 20 FT CONTOUR
- 100 FT CONTOUR
- 2500 FT CONTOUR
- GATE
- CAMP
- PAVED PUBLIC ROAD
- GRAVEL PUBLIC ROAD
- YEAR ROUND GRAVEL
- SUMMER ROAD (NO/LIMITED GRAVEL)
- IMPROVED WINTER ROAD
- UNIMPROVED WINTER ROAD
- SKID TRAIL/UNMAINTAINED ROADWAY
- UNKNOWN ROAD SURFACE
- CUT/RU EDGE


Map Document: (D:\map_projects\VT\clients\PLUM_CREEK_7086\HARVEST_UNITS\LM-03-01-09_CLOUGH_BROOK_NORTH.mxd)
10/21/09 -- 11:27:20 AM

568000 568500 569000 569500 570000 570500

568000 VT STATE PLANE NAD 83 METERS 568500 569000 569500 570000 570500

**LAND OF PLUM CREEK MAINE
TIMBERLANDS, LLC.
NORTHERN KINGDOM UNIT-VT
HARVEST MAP
CLOGH BROOK NORTH
LM-03-01-09**

SCALE 1:15,840 1"=1,320'
0 660 1,320 2,640 Feet



Town: LEMINGTON
County: ESSEX
Compartment: LM-03
Forester: JCH
Job Number: LM-03-01-09
Tractor: TBD

STAND_NUMB	STRATA	OS_TYPE	SILV_PRES	ACRES
LM-03-24	SH34B/SH2B	SH3B	ITH	68.8
LM-03-34.0	H3B	H3B	2SS	137.0
LM-03-43.0	HS3B	HS3B	OSR/ITH	115.1
LM-03-44	H3B	H3A	ITH	37.0
LM-03-46	H3C/H1B	H3C	OSR	15.9
LM-03-54	H3B	H3B	STC	97.4
TOTAL (Acres derived from GIS)				471.1

- PROPOSED HARVEST
- ACTIVE HARVEST
- COMPLETE HARVEST
- APPROVED HARVEST
- POSTPONED HARVEST
- FLOWAGE/BOG
- GRAVEL PIT
- LANDING
- OPEN/AGRICULTURAL
- RIPARIAN ZONE
- SPECIAL TREATMENT AREA
- WATER
- COUNTY LINE
- TOWN LINE
- STREAM
- STAND LINE
- COMPARTMENT LINE
- GATE
- CAMP
- PAVED PUBLIC ROAD
- GRAVEL PUBLIC ROAD
- YEAR ROUND GRAVEL
- SUMMER ROAD (NO/LIMITED GRAVEL)
- IMPROVED WINTER ROAD
- UNIMPROVED WINTER ROAD
- SKID TRAIL/MAINTAINED ROADWAY
- UNKNOWN ROAD SURFACE
- CUT/HU EDGE
- STAKE AND STONE PILE

Map Document: (D:\map_projects\VT\clients\PLUM_CREEK_7086\HARVEST_UNITS\LM-03-01-09_CLOGH_BROOK_NORTH.mxd)
10/21/09 -- 11:27:20 AM

*PHOTO TAKEN 2006

Exhibit 2.

“Silvicultural Guide for Northern Hardwood Types in the Northeast (revised)” by Dr. William B. Leak, et al., published by the United States Department of Agriculture, Research Paper NE-603 (the “Guide”).



United States
Department of
Agriculture

Forest Service

Northeastern Forest
Experiment Station

Research Paper NE-603



Silvicultural Guide for Northern Hardwood Types in the Northeast (revised)

William B. Leak
Dale S. Solomon
Paul S. DeBald

Exhibit
2.



Abstract

A revision of the 1969 silvicultural guide for northern hardwoods, provides up-to-date information on both even-age and uneven-age silviculture and management for beech-birch-maple, beech-red maple, and mixedwood stands in the Northeast.

The Authors

WILLIAM B. LEAK is principal silviculturist with the Northeastern Forest Experiment Station at Durham, New Hampshire. DALE S. SOLOMON is principal mensurationist with the Northeastern Station at Orono, Maine. PAUL S. DEBALD is a research forester with the Northeastern Station at Delaware, OH.

Manuscript received for publication 24 September 1986.

Northeastern Forest Experiment Station
370 Reed Road, Broomall, PA 19008
August 1987

COVER—Many-aged northern hardwood stand after improvement cutting, Bartlett Experimental Forest

Purpose and Scope

This is the third silvicultural guide for northern hardwoods (beech-birch-maple) in the Northeast. The first, published in 1958 (Gilbert and Jensen 1958), provided general guidelines on initial cutting methods in uneven-aged old-growth stands and even-aged second-growth stands. The second, published in 1969 (Leak et al. 1969), provided quantitative information on stocking and yield as well as a key to specific stand prescriptions, particularly for even-age management. This revised guide includes new information on forest types, site, stocking, growth and yield, and regeneration methods, including shelterwood and group selection.

The information in this guide applies to about 20 million acres of northern hardwood and mixed hardwood-conifer types across New England and New York. Outside this area, the guide should be applied with caution. The guide is primarily concerned with timber production. A guide to the management of wildlife habitat in hardwood and conifer types in New England is in preparation.

Regional Conditions

Northern hardwoods and associated species are used for a variety of products, including veneer, sawlogs, boltwood, pulpwood, fuelwood, and miscellaneous products such as posts. In New England, sawlogs and veneer presently account for about one-quarter of the hardwood harvest, and pulpwood accounts for about one-half. In states such as New Hampshire, fuelwood currently accounts for a significant proportion of the hardwood cut. In many areas, markets for low-quality material provide opportunities for improving the northern hardwood forest without heavy investments in noncommercial silvicultural work. Also, substantial increases in land and timber prices over the last 10 to 15 years have strengthened timberland investments. Indications are that current trends will continue upward, though perhaps at a lower rate.

Many timberland owners in New England own land primarily for reasons other than growing timber. In New Hampshire and Vermont, only 6 percent of the owners, controlling 21 percent of the timberland, listed wood production as one of the important reasons for ownership (Kingsley and Birch 1977). Foresters must consider the other values of timberland—recreation, esthetics, buffers, wildlife, investment, etc.—and be prepared to use silvicultural approaches that will complement or enhance these values.

Species and Sites

Three cover types, or subtypes, are described in this guide: beech-birch-maple (typical northern hardwood), beech-red maple, and mixedwood (hardwoods and associated softwoods). The occurrence of these types usually is related to site conditions—soils, climate, and bedrock mineralogy; in parts of the Northeast, these forest types are known to occur on certain land types, forest habitats, or soil series. Those who manage stands primarily for paper birch, oak, white pine, spruce-fir, or cherry-maple should use the guides written specifically for those types.

The beech-birch-maple type contains sugar maple as the characteristic species in proportions ranging from 15 to 20 percent to nearly 100 percent of the basal area. This type is characteristic of well- to moderately well-drained, fine-textured or loamy till soils. Sugar maple and/or white ash are most abundant on the best soils—for example, those that are enriched with organic matter or derived from limestone. However, on average beech-birch-maple sites, beech may account for up to 50 percent of the basal area. The most common birch species are yellow and paper birches. However, in southern New England, sweet birch and northern red oak (often of good quality) may be common associated species. The successional tendency is toward the tolerant species—beech and sugar maple.

Beech-red maple stands usually occupy poorer sites than beech-birch-maple stands—soils that are more shallow, wetter, or drier than those with typical northern hardwoods. The central characteristic of these hardwood stands is that sugar maple is uncommon and/or slow growing. On dry sites, beech may be the predominant species. On wet sites or shallow soils, red maple often is the most common species. Yellow birch and paper birch (or sweet birch in southern New England) are common associates. Some of these stands originated from heavy cutting of softwood or mixedwood stands. Old stands sometimes show a successional trend toward tolerant softwood types—hemlock and/or red spruce. Or the successional tendency may be toward a predominance of beech. Some of the characteristic soils are productive for red oak; less so for sugar maple, yellow birch, or white ash.

Mixedwood (hardwoods with primarily spruce, hemlock, or balsam-fir) stands have at least three origins: (1) partial or heavy cutting of softwood stands, which allowed hardwood invasion to occur; these stands appear as the beech-red maple cover type with a softwood component or understory; (2) forest succession on abandoned fields, which tends to favor increased proportions of white pine, hemlock, spruce (red or white), or balsam-fir even on good hardwood sites that normally would support the beech-birch-maple type; and (3) diverse site conditions, e.g., shallow or rocky soils interspersed with better soils, which results in groups of hardwoods and softwoods. Because of different origins, successional trends in mixedwood stands may be toward tolerant softwoods, hardwoods, or maintenance of the mixedwood character.

Type	Distinguishing Factors
Beech-birch-maple	<ol style="list-style-type: none"> 1. Fifteen percent or more sugar maple in overstory and/or understory. 2. Sugar maple more abundant than red maple. 3. Less than 25 percent softwoods in overstory or understory. 4. Commonly found on well- to moderately well-drained fine-textured till soils.
Beech-red maple	<ol style="list-style-type: none"> 1. Less than 15 percent sugar maple in overstory and understory. 2. Red maple more abundant than sugar maple. 3. Less than 25 percent softwoods in overstory and understory. 4. Commonly found on soils that are wetter, drier, more shallow to bedrock or pan, rockier, or more poorly aerated than those supporting beech-birch-maple stands.
Mixedwood	<ol style="list-style-type: none"> 1. Softwoods (primarily spruce, hemlock, or balsam-fir) account for 25 to about 65 percent of the overstory and/or understory. 2. No specifications on the relative proportions of sugar maple and red maple. 3. Found on a wide variety of soils due to the varied past history and origin of mixed-wood stands.

Note that certain intolerant or intermediate species—paper birch, yellow birch, aspen, etc.—are not indicative of site condition.

Silvicultural opportunities differ for each subtype. The beech-birch-maple type regenerates and produces large, high-quality hardwoods such as sugar maple, yellow birch, or, in some areas, white ash. The beech-red maple type commonly produces smaller, lower grade sawlogs. However, such areas can produce acceptable boltwood stems (birches) or aspen for wildlife; oak sawlogs are a possibility in certain parts of the northern hardwood region. In many mixedwood stands, the possibilities for hardwood products are similar to those for the beech-red maple subtype. Here, volume production can be increased by gradually favoring the softwood component. However, due to the variable origin of mixedwood stands, large high-quality hardwoods can be grown on some sites.

Silvical characteristics of the important species found in these three types are outlined in Table 1. The shade-tolerance categories of tolerant, intermediate, and intolerant indicate whether the species can regenerate and persist under conditions of heavy, moderate, or no shade, respectively. Moderate shade would be about 30 to 70 percent crown cover.

The categories of early height growth indicate the general ability of a species to outgrow its associates up through the sapling stage with no overhead competition. For example, aspen grows very fast and seldom needs release from associated species to maintain a dominant position, and red maple normally is outgrown by aspen and paper birch. Sugar maple and beech are outgrown by most other hardwood species even when free of competition.

The relative site requirements vary by species. White ash, and sugar maple to a slightly lesser extent, are most abundant and reach best development on the best soils. Conversely, spruce, hemlock, and white pine are common and grow to fairly large size on soils that are shallow, rocky, sandy, or wet. Paper birch, red maple and aspen, though fast growers on good sites, can be reproduced and grown fairly successfully on poor sites.

The natural pruning categories refer to a species' relative ability to produce clear boles. The softwoods, as well as sugar maple and beech, only produce clear boles at an early age under conditions of high stand density. However, paper birch, white ash, and aspen produce clear straight boles at only 50 to 60 percent of maximum stand density.

Protection

Diseases and insects are the two important problems. Fire damage and prevention have a minor impact on silvicultural methods in northern hardwoods and related types.

Table 1.—Silvical characteristics of the important species of the three cover types

Species	Shade tolerance	Early relative height growth	Relative site requirements	Natural pruning	No. years between good seed crops	Sprouting vigor	Delayed germination
Sugar maple	Tolerant	Slow to moderate	High	Poor to medium	3-7	Moderate—small stumps	Negligible
American beech	Tolerant	Slow	Medium	Poor	2-5	Low—stump sprouts High—root suckers	None known
Yellow birch	Intermediate	Moderate	Medium to high	Medium	1-3	Low	Seldom
Paper birch	Intolerant	Fast	Medium	Good	2	Moderate—small stumps	None known
White ash	Intermediate (more tolerant as a seedling)	Moderate	High	Good	2-5	Moderate to high	Up to 75 percent
Red maple	Intermediate	Moderate	Low	Medium	1	High	Moderate percentage may germinate 2nd spring
Aspen	Intolerant	Very fast	Low	Good	4-5	High—root suckers	None
Northern red oak	Intermediate	Moderate	Medium	Medium	3-5	High	None
Red spruce	Tolerant	Very slow	Low	Poor	3-8	None	None known
Eastern hemlock	Tolerant	Very slow	Low	Poor	2-4	None	None known
Eastern white pine	Intermediate	Slow to moderate	Low	Poor	3-10	None	None known

Few silviculturalists can develop expertise in entomology or pathology. However, they should be able to: (1) recognize threats to quality and vigor; (2) apply silvicultural and utilization methods that will minimize losses; and (3) know when to seek expert advice on potential outbreaks. Beyond this section, few references will be made to specific insects and diseases. However, many of the silvicultural practices described later relate to potential stand quality and risk.

Microorganisms that cause decayed and discolored wood, one of the most widespread problems in both hardwoods and softwoods, enter through dying or damaged roots, stems, or branch stubs. Other fungi cause collar cracks in birches and root rot and basal decay in many other species. For details on discoloration, decay, and other defects, see Shigo and Larson (1969) and Shigo (1983).

To limit losses in stand volume and quality from decay and discoloration, silvicultural methods should be directed toward (1) removing defective trees during harvest operations, (2) encouraging the development of small limbs and early natural pruning by maintaining fairly high stand densities in sapling and pole stands (or consider artificial pruning), and (3) minimizing logging damage to roots, stems, and branches. Decay and discoloration associated with wounds or branch stubs usually do not enter into wood formed after the wound occurs or after the branch dies.

Nectria cankers are common in many hardwoods, and *Eutypella* canker is found on maples. Associated decay is limited, but wind breakage is common and the cankers often cause quality defects in the most valuable parts of the bole.

Beech-bark disease is the most lethal disease of beech. The beech scale (*Cryptococcus fagi* (Baer)) punctures the bark, allowing a bark-killing fungus (*Nectria coccinea* vac. *faginata*) to enter. Small vigorous trees sometimes survive the disease, though quality may be severely reduced. Large trees seldom survive. Significant tree-to-tree variation is evident in susceptibility to both the scale and the *Nectria* fungus. Trees exhibiting the characteristic red fruiting bodies should be harvested without delay.

The saddled prominent (*Heterocampa guttivitta*) is the most serious defoliator of beech-birch-maple stands. White ash and red maple sometimes escape attack, but most other hardwoods can be heavily defoliated. The outbreaks occur on about a 10-year cycle, and last 1 to 3 years. Defoliation in 2 successive years may be followed by widespread mortality and growth losses, as well as degrade (e.g., stain) from secondary insects and organisms. Aerial chemical sprays have proven effective in limiting damage by the saddled prominent.

Gypsy moth (*Lymantria dispar*) is the most serious defoliator of oak in central to southern New England. Grey birch, aspen, and sometimes other hardwoods are frequently defoliated as well. White pine and hemlock associated with oak stands also are defoliated during severe attacks. Outbreaks generally last 1 to 3 years. Mortality and/or growth loss in oak can be significant after two or more defoliations. There is some indication that degrade from insects (e.g., oak borer) may increase in trees weakened by defoliation. Insecticidal and biological aerial sprays are effective.

Larval galleries of the sugar maple borer (*Glycobius speciosus*) result in partial girdles and cankers that allow the development of decay and discoloration and increase susceptibility to wind breakage. Up to 50 to 60 percent of the sugar maple in a stand may be affected, with an average of two to three significant cankers per tree. Contrary to earlier recommendations, exposure of trees by cutting apparently does not increase susceptibility to damage by the sugar maple borer.

Management Objectives and Approaches

In developing a long-range timber management objective for an entire forest property, the owner or manager should first decide on product and species goals. A common product objective is high-quality material for veneer logs, sawlogs, and boltwood combined with utilization of the poorer quality material for pulp or fuelwood. A second objective on certain industrial lands or private ownerships might be maximum production of wood for fiber and/or fuel. Other product objectives include high-intensity management for veneer logs, or production of small logs and boltwood. The choice of species or species group is closely related to the desired product, and is concerned with what species to favor in improving and regenerating the stand.

Two primary factors affecting the choice of product and species are (1) current and projected markets, and (2) the capability of the land to grow certain products and species. Once the product and species objectives are set, a preliminary decision can be made on appropriate silvicultural systems for the property. For tolerant species and high-quality products, one standard approach in the Northeast has been uneven-age management using individual-tree selection or some form of partial cutting. For quality products from intolerant and intermediate species, a common approach is even-age management using clearcutting and intermediate treatments. Even-age management with clearcutting is best suited to the mechanized production of fiber or fuel.

Table 2.—Species of regeneration favored (not exclusively) by certain harvesting methods in three cover types

Type	Individual tree-selection	Group selection	Dense shelterwood ^a	Open shelterwood ^b	Clearcut
Beech-birch-maple	Sugar maple Beech	Birches	Sugar maple Beech	Yellow birch	Birches
Beech-red maple	Beech Red maple	Red maple Birches	Beech Red maple	Red maple Yellow birch	Birches
Mixedwood	Tolerant softwoods or hardwoods	Red maple Birches	Hemlock Spruce Tolerant hardwoods	Red maple Birches	Birches

^aResidual crown cover of about 80 percent.

^bResidual crown cover of 30 to 50 percent (occasionally up to 70 percent).

However, in assessing each stand, additional factors must be considered in reaching a decision on the immediate silvicultural techniques. Either an uneven-age or even-age approach can be used to grow most species groups or products: uneven-age management with group selection can ensure a good mix of intermediate and some intolerant species; even-age management with shelterwood cutting can be designed to encourage a high proportion of tolerants (Table 2). And rotation age, stocking, stand structure, and logging equipment can be varied to meet various product objectives.

One of the important additional factors to consider is current stand condition. For example: high-risk stands may need to be regenerated by clearcutting or shelterwood to prevent large volume losses; or clearcutting a stand with a wide range in tree diameter may remove a high proportion of financially or biologically immature trees.

Another factor is accessibility. With high costs for road construction, some form of heavy cutting may be the only economically feasible regeneration harvesting method on the first entry. Esthetic and wildlife objectives also should be considered in choosing a silvicultural system. And in special circumstances, the possibility of site and/or stand deterioration needs to be assessed. For example, clearcutting on very poorly drained soils without adequate advance regeneration or in potential frost pockets may result in an overabundance of herbaceous or shrubby vegetation.

Uneven-Age Management

Harvesting Methods

Uneven-age management is implemented by individual-tree selection and group selection. Individual-tree selection removes trees one by one to maintain a fairly uniform and continuous crown cover appropriate for regenerating tolerant species. Group selection is the removal of trees in groups roughly 1/20 to 2 acres in size. It is especially appropriate where: (1) the objective is to maintain up to one-half of the regeneration in intolerant or intermediate species, and (2) the overstory contains groups of poor-risk, defective, or overmature trees. Group selection generally is applied in combination with individual-tree marking between the groups.

Group selection may be applied in two ways: groups of overstory trees can be removed, leaving a desirable stand of seedlings, saplings, or small poles; or entire groups of trees down to 2-inches d.b.h can be removed. The latter approach is used to eliminate undesirable sapings and small poles, resulting in a maximum proportion of intolerant or intermediate regeneration.

Growth and Yield

Results from a study of residual basal area and structure in a second-growth, beech-red maple stand illustrate the typical responses of hardwood stands in New England to density and structure (Table 3). Basal-area and cubic-foot

Table 3.—Net annual growth per acre over a 9-year period of sawtimber (10.5 inches plus) and total stand (4.5 inches plus), by residual basal area and amount of sawtimber in a second-growth beech-red-maple stand (approximate site index 55 for sugar maple) (Solomon 1977)

Residual basal area (ft ²)	Residual sawtimber		Net growth				Accretion	
	Percent	Basal area	Sawtimber ^a		Total		Sawtimber ^b	
			Square feet	Board feet	Square feet	Cubic feet	Square feet	Board feet
40	30	12	1.4	126	2.2	53	.51	46
	45	18	1.3	117	2.0	48	.68	61
	60	24	0.5	45	1.8	43	.69	62
60	30	18	1.3	117	1.7	41	.55	50
	45	27	1.7	153	2.3	55	.76	68
	60	36	1.6	144	2.0	48	.88	79
80	30	24	2.0	180	1.7	41	.64	58
	45	36	1.6	144	1.7	41	.71	64
	60	48	1.0	90	1.2	29	.92	83
100	30	30	1.9	171	1.7	41	.70	63
	45	45	1.6	144	1.2	29	.88	79
	60	60	1.1	99	0.9	22	1.07	96

^aIncludes only the sawtimber portion of the stem.

^bBased on conversions of 90 board feet and 24 ft³ per square foot of basal area.

growth generally are best between 40 and 60 ft² of residual basal area per acre. Board-foot growth, however, is best between 60 and 80 feet of basal area provided that at least 25 to 35 ft² of basal area in sawtimber is maintained. With these fairly low sawtimber basal areas, much of the sawtimber growth is ingrowth into the larger sizes rather than accretion. Accretion tends to rise as the basal area in sawtimber approaches 50 to 60 ft²; however, mortality and lower sawtimber ingrowth reduce the net sawtimber growth. Retention of live branches is noticeable at 60 ft² basal area, indicating that timber quality development should be best at residual basal areas of 70 to 80 ft².

On better sites, those supporting beech-birch-maple stands, experience indicates that higher amounts of residual sawtimber could be maintained that produce high-quality sawtimber growth.

Stocking and Structure

Recommended minimum residual basal areas in trees 5.0 inches and larger in d.b.h. are:

Type	Residual Basal Area (ft ²)
Beech-birch-maple	70 (65–75)
Beech-red maple	70 (65–75)
Mixedwood	100 (80–120)

The mixedwood goal applies to residual stands with 25 to 65 percent softwood in trees 5.0 inches and larger in diameter. A range in basal area is given to encourage flexibility. On good sites, those with an abundance of quality timber, residual basal areas above the suggested range may be specified. However, residual basal areas below the suggested range may impair quality development.

Stand structural goals for uneven-age management are specified by the maximum size tree to leave, and the diameter distribution. Diameter distributions are approximated by a reverse J-shaped curve, with a slope defined by *q*—the quotient between numbers of trees in successively smaller d.b.h. classes.

The structural goals in this guide are based on a range in *q* from 1.3 to 1.7, and a maximum tree size (for general planning purposes) of 20 inches d.b.h.:

<i>q</i>	Approximate percent sawtimber
1.7	45
1.5	55
1.3	70

We emphasize that the maximum tree size of 20 inches d.b.h. is a very flexible goal. Tree vigor and quality are more important than the specified maximum tree size in deciding which trees to take or leave. On poor sites, tree vigor and quality of some species may decline rapidly at 16 inches d.b.h. or larger; on these sites, low amounts of sawtimber ($q = 1.7$) are most appropriate. On good sites, trees may easily be grown to 24 inches or larger; on such sites, high proportions of sawtimber ($q = 1.5$ to 1.3) should be best.

On the basis of these combinations of q and residual basal area, residual structural goals in terms of basal area per acre by diameter class are outlined in Table 4 for both hardwood and mixedwood types. Only three diameter classes are used since this results in easier application and allows for some departure from the strict reverse J-shaped form. Recent information indicates that slightly S-shaped form of diameter distribution may be more natural, productive, and economical.

In choosing a structural goal, it often is reasonable to aim for a q that is about the same as or slightly lower than the existing q before cutting. The q before cutting can be judged quickly by using the tabulation for percent sawtimber. A more precise estimate of the appropriate residual structure can be developed by following the marking guide procedures described in the next section. The structures listed in Table 4 should be used as a guide, and can be attained by feasible and economical cutting practices.

The structural goal of $q = 1.7$ is appropriate for the first entry in many cutover stands, which often have a low proportion of sawtimber (Fig. 1). However, the initial cut in a previously unmanaged stand may produce an extremely variable diameter distribution. Total residual density, and the removal of poor growing stock, are more important than structure in these early cuts. During subsequent entries, it may be feasible to leave a higher proportion of sawtimber (a lower q). On mediocre sites (e.g. beech/red maple stands), the sawtimber will decline in vigor and growth rate as it becomes larger, so it may never be possible to grow large-size trees or to reduce the q below 1.5 or 1.7. On good sites (e.g., supporting sugar maple/ash) capable of sustaining high proportions of sawtimber (Fig. 2), q 's of 1.3 are attainable.

On areas scheduled for maximum production of fiber or fuel, low proportions of sawtimber (a q of 1.7 or higher) should be best.

In poletimber stands with less than about 25 to 30 ft^2 of sawtimber, there is little reason to be concerned about structure. Such stands can be treated by commercial stand improvement measures that remove the poorer quality overstory stems and leave 70 to 80 ft^2 of basal area per acre. In subsequent treatments, as the sawtimber component develops, the use of structural goals will be more appropriate.

Table 4.—Minimum stand structure objectives for residual hardwood (beech-birch-maple and beech-red-maple) and mixedwood stands

D.b.h. class (inches)	$q = 1.7$		$q = 1.5$		$q = 1.3$	
	Hardwood	Mixed-wood ^a	Hardwood	Mixed-wood ^a	Hardwood	Mixed-wood ^a
	----- ft^2 of basal area/acre-----					
6-10	38	54	30	42	21	30
12-14	18	26	20	28	20	28
16+	14	20	20	30	29	42
All	70	100	70	100	70	100

^aSoftwood basal area 25 to 65 percent of total.



Figure 1.—Cutover stand of northern hardwoods with low to moderate proportion of sawtimber. Immediate residual goals in such stands would be 30 to 40 ft² of sawtimber or a q or 1.7 to 1.5. A good northern hardwood site with good-quality sugar maple, this area could support more sawtimber (lower q) in the future.

Cutting Cycle

The cutting cycle—the years between harvests on the same area—is determined by accessibility, the need for an economic harvest, timber growth, and quality. Based on an average basal growth rate for northern hardwoods of 2 ft²/acre/year a residual stand of 70 ft² will grow to 100 ft²—a reasonable maximum—in about 15 years. At that time, a cut of about 30 ft² (8 to 9 cords) will be available. With good accessibility, high timber values, or high risk of damage from insects or diseases, the cutting cycle might be lowered to 10 years. Under opposite conditions, cutting cycles of up to 20 years sometimes are used. With long cutting cycles, the total residual basal areas in Table 4 can be lowered to minimums of 60 to 65 ft² for hardwood and about 75 to 80 ft² for mixedwood, proportioning the reductions among size classes.

Marking Guides

The success of uneven-age management—both silviculturally and economically—depends to a large extent on the choice and application of appropriate marking guides.

Marking guides can be developed from a prism inventory of the stand, classifying the trees by d.b.h. and condition classes:

1. Acceptable Growing Stock: Trees with the potential to produce sawlog or better quality material, now or in the future, and that are in vigorous condition. These can be subdivided into mature (based on biological or financial maturity) and immature. Suggested tree sizes denoting financial maturity are shown in Table 5.



Figure 2.—Northern hardwood stand with a high proportion of fairly good-quality sugar maple. The residual goal in this stand would be about 50 ft² of sawtimber or a q of approximately 1.3.

2. Unacceptable Trees: Trees that will not produce sawlog or better material now or in the future or trees that are high risk—subject to mortality or rapid losses of merchantable volume or quality before the next harvest. Valuable high-risk trees are especially important to recognize.

3. Cull: Trees with more than 50 percent of their cubic volume in sound or rotten cull; or use a local or agency definition.

Certainly, additional tree condition classes could be developed to meet local timber or wildlife needs. Individual species or species groups often should be tallied to help refine silvicultural objectives and develop marking guides. Prism-plot basal areas, by d.b.h. and tree condition class can be summarized as in the following example:

D.b.h. class (inches)	-ft ² basal area/acre-					Total
	Mature	Imma- ture	Defec- tive	High risk	Cull	
6-10	—	40	10	—	—	50
12-14	—	10	10	20	—	40
16+	5	5	5	10	—	25
All	5	55	25	30	—	115

Table 5.—Tree-size objectives (d.b.h.) denoting peak of possible log-grade improvement (financial maturity) for northern hardwoods (high and medium-grade potential reflect the marker's best judgement on tree condition and site potential for that species)

Species	Grade potential ^a	Log section ^b	D.b.h. objective
			<i>Inches</i>
Yellow birch	High	1	18
	High	2	20
	Medium	1	16
	Medium	2	18
Sugar and red maple ^c	High	1	18
	High	2	20
	Medium	1	16
	Medium	2	18
Beech	High	1	16
	High	2	18
	Medium	1	14
Paper birch	Medium	2	16
	High	1	16
	High	2	16
	Medium	1	12
White ash	Medium	2	12
	High	1	18
	High	2	20
Red spruce	Medium	1	16
	Medium	2	18
	Medium or better	Any	14-16
Hemlock	Medium or better	Any	18-20

^aHigh-grade potential means that the first 1- or 2-log portion of the stem could produce grade 1 or veneer-grade logs. Medium-grade potential means that the highest quality would be no more than grade 2. Grades are based on USDA Forest Service standard specifications for hardwood factory lumber and veneer logs.

^bPredominant merchantable height.

^cIn many areas, red maple has medium-quality potential at best, so d.b.h. objectives commonly are 14 to 16 inches.

This hypothetical northern hardwood stand has 57 percent of the basal area in sawtimber. A stand with a q of 1.5 has approximately 55 percent of the basal area in sawtimber, so a structural goal of $q = 1.5$ would be a reasonable first choice (see tabulation for percent sawtimber). However, in looking at the tree condition classes, a reasonable set of marking guides might be:

1. Remove high-risk timber in the 12- to 14-inch class.
2. Remove high-risk timber in the 16+ class.
3. If a cordwood market were available, remove defective trees in the 6- to 10-inch class

Application of these marking rules would leave a residual stand as follows:

D.b.h. class (inches)	Mature	Immature	Defective	Total	Goal ($q = 1.7$)
-----ft ² of basal area/per acre-----					
6-10	—	40	—	40	38
12-14	—	10	10	20	18
16+	5	5	5	15	14
All	5	55	15	75	70

This residual stand is close to the structural goal of $q = 1.7$ in Table 4, and the stand contains a high proportion of vigorous growing stock.

An alternative approach is to follow generalized marking rules related to stand condition (Table 6). With 65 ft² of sawtimber, the hypothetical initial stand qualifies as beech-birch-maple large sawtimber. Total initial basal area is more than 100 ft², so a cut is warranted using either single-tree or group selection. The suggested residual basal area is 75 ft², with 40 to 55 ft² of sawtimber. In this example, the use of Table 6 leads us into leaving a little more sawtimber basal area (8 to 23 ft² more), and a corresponding smaller amount of poletimber, than is suggested by the prism-plot summary of the actual condition of the growing stock.

Noncommercial Work

Noncommercial stand improvement work is minimal with uneven-age management because each harvest operation provides the opportunity to improve the stand by removing marginal or submarginal trees. However, three types of noncommercial work are possible.

Table 6. Summary of general marking guides

Type	Stand size class	Initial stand		Cutting method	Regeneration favored	Residual stand	
		Sawtimber basal area	Total basal area 5.0 inches + ^a			Sawtimber basal area	Total basal area 5.0 inches + ^a
Beech-birch-maple	Poletimber	30	100 +	Stand Improvement	Sugar maple-beech	65	65
	Small sawtimber	30-50	100 +	Single-tree Group	Sugar maple-beech Yellow birch-paper birch	30-40 30-40	70 70
	Large sawtimber	50-75	100 +	Single-tree Group	Sugar maple-beech Yellow birch-paper birch	40-55 40-55	75 75
Beech-red-maple	Poletimber	30	100 +	Stand Improvement	Beech-red maple	65	65
	Small sawtimber	30-50	100 +	Single-tree Group	Beech-red maple Yellow birch-paper birch	30-40 30-40	70 70
	Large sawtimber	50-75	100 +	Single-tree Group	Beech-red maple Yellow birch-paper birch	40-55 40-55	75 75
Mixedwood	Poletimber	50	130 +	Stand Improvement	Tolerant softwoods or hardwoods	80	80
	Small sawtimber	50-70	130 +	Single-tree Group	Tolerant softwoods or hardwoods Intermediate ^c and intolerant hardwoods	45-55	100
	Large sawtimber	70-90	130 +	Single-tree Group	Tolerant softwoods or hardwoods Intermediate ^c and intolerant hardwoods	55-75	100-120
				Group	55-75	100-120	100-120

^aStands with less basal area than specified should be left to grow.

^bGroup selection normally includes individual-tree selection between groups.

^cIf tolerant or intermediate softwood regeneration is present, group selection can be used to favor these species by group removal of the overstory where the softwood regeneration is well developed.

1. **Cull Removal:** Culls to be removed should be designated during marking. Residual culls should be included in defining the residual basal area and structure—because they influence growth, regeneration, and quality development. Elimination of understory culls is of doubtful value because of their abundance, slow growth, and high mortality. Keep in mind the importance of reserving some cull trees, especially the large ones, for wildlife. More than 30 species of birds and 20 mammals use culls in New England for feeding, nesting, or denning.
2. **Two-Inch Removal:** When group selection is applied, and the saplings or small poles within a group are of undesirable species or quality, complete stem removal down to about 2 inches d.b.h. is recommended. Generally, this is most easily done during the logging operation, though a separate postlogging treatment also is possible using chemical or mechanical methods.
3. **Group Stand Improvement:** Groups of saplings or small poles resulting from previous group-selection cuttings may be dominated by stems of low-quality or undesirable species. If less than about 40 percent of the overstory stems are minimally acceptable, consider a noncommercial weeding/cleaning to be carried out in the substandard groups. This work is expensive and should be restricted to those instances where the site is good and a marked increase in stand value will result. The operation should be the minimum required to create an overstory with at least 40 to 50 percent of the stems in acceptable species and quality.

Regeneration

Under single-tree selection, more than 90 percent of the regeneration will be tolerant species. Under group selection, based on groups averaging about one-half acre in size with all stems larger than 2 inches removed, about one-third of the regeneration will be of intermediate or intolerant species (Table 7). In larger groups up to about 2 acres in size, the intolerant and intermediate species should account for one-half of the regeneration. Limited experience indicates that the proportion of intolerants (e.g., paper birch) may equal the proportion of intermediates in groups two-thirds of an acre or larger in size.

Species composition of the regeneration under single-tree selection is closely related to the advance regeneration. Areas to be regenerated to sugar maple or softwoods should show evidence of these species in the existing advance regeneration. To be considered well established, a hardwood or hemlock stem should be 3 to 4 feet tall; a spruce or fir, about 1 foot tall.

Under group selection, the presence of residual tolerant stems in the seedling, sapling, or small-pole sizes will limit

Table 7.—Species composition of stocked mil-acres, 10 to 15 years after cutting in beech-birch-maple stands, by tolerance group and cutting method

Tolerance group ^a	Clearcutting	Group selection	
		Group selection	Single-tree selection
-----Percent-----			
Tolerant	43	62	92
Intermediate	19	34	7
Intolerant	38	4	1

^aTolerant: beech, sugar maple, eastern hemlock, and red spruce (also balsam-fir if present); intermediate: yellow birch, white ash, and red maple; intolerant: paper birch and aspen.

intolerant or intermediate regeneration. In regenerating birch under group selection, advanced regeneration larger than 2 inches d.b.h. should be sparse, or it should be removed in creating the opening (Fig. 3). Snow-free logging generally is more effective than winter logging in reducing unwanted advanced growth.

Group openings in poletimber stands do not always regenerate well—especially on sites that are extremely wet or dry—and are not recommended. Regeneration in groups receives more snow damage than in clearcuttings, apparently due to the extra snow that drifts into small canopy openings. But research has indicated that less than 10 percent of the groups in hardwood and mixedwood stands were in poor condition due to snow damage.

Border-tree quality is a consideration in group selection because border trees may tend to retain live limbs or produce epicormic sprouts. Poles or small sawtimber of hardwood species should not be left as border trees without trainers or buffer trees if they have (1) clear boles but small live crowns; or (2) lower live limbs that will seriously detract from quality if allowed to persist. White ash and paper birch are least likely to produce epicormic sprouts, but live limbs on any hardwood or softwood species will remain alive for a substantial period of time if exposed to full sunlight.

What constitutes adequate stocking in the seedling and sapling size classes of an uneven-aged stand? This question has not been answered completely for New England hardwood and mixedwood stands. However, in extensively



Figure 3.—Portion of a group-selection opening in nearly mature northern hardwoods. Complete removal of the understory will result in maximum amounts of intolerant-intermediate regeneration.

managed stands, the percentage of milacres stocked with at least one stem between 3 feet tall and 1.5 inches d.b.h. usually exceeds 65 percent. Percent stocking of desirable species much lower than this—below 50 percent, for example—would indicate the need for special attention to regeneration, perhaps the use of small group-selection openings. The number of stems of commercial species (1.5 to 4.5 inches, or in the 2, 3, and 4-inch classes) commonly ranges from about 200 to 450. If adequate stocking in the seedling class is present, but 2- to 4-inch saplings seem deficient, a harvest cutting to the recommended residual basal area should solve the problem. There is no concensus at present on the need for mechanical or chemical treatments to improve the composition of the seedling-sapling component under single-tree selection. Work on the Bartlett Experimental Forest with single-stem timber stand improvement in understory beech produced little permanent change at high cost. However, the subject deserves further study.

Even-Age Management

Harvesting Methods

Two even-age harvest cutting methods commonly used in the Northeast are clearcutting and shelterwood. The seed-tree method also is recommended sometimes for large cutting areas where the available seed source of desired species is limited.

Clearcutting is the harvesting of all merchantable trees on an area generally followed by a chemical or mechanical removal of trees down to 2 inches d.b.h. (Fig. 4). Sometimes groups of trees larger than 2 inches d.b.h. are left if they are of desirable species. Isolated residual trees may develop large limbs and poor quality. However, pole-size or larger sugar maple with good crowns, clear boles, and no tendency to produce epicormic sprouts will experience little



Figure 4.—Clearcutting in northern hardwoods with essentially complete removal of the understory.

degrade when left as residuals to increase the tolerant component of the new stand. Clearcutting boundaries can be designed to follow natural stand or topographic boundaries to minimize adverse esthetic impacts.

Progressive stripcutting is a variation of clearcutting that is especially well suited to the regeneration of yellow birch and other intermediate species. Strips 50 to 100 feet wide are laid out along the contour. In the first cutting, every third strip is removed; 2 to 4 years later, one strip next to each initially cut strip is removed. After another 2- to 4-year interval, the final strips are cut. Trees down to 2 inches d.b.h. are removed or felled. The material generally is skidded down the strips currently being cut, which results in a high degree of scarification and the removal of most undergrowth.

In mixedwood stands on wet areas, where windthrow would be a threat under unevenage or shelterwood systems, narrow strips (30 to 50 feet wide) sometimes are used to help perpetuate a softwood component. Winter logging usually is advisable in these areas, and softwood regeneration is most probable if the strips contain established softwood seedlings.

In the Northeast, the shelterwood system commonly is applied in two cuts: an initial seed cut (Fig. 5) and a final removal cut. Where the objective is to regenerate tolerant species and little or no advanced regeneration is present, an initial light preparatory cut also may be desirable to stimulate seed production of desirable species and the establishment of small seedlings. For tolerant regeneration, the seed cut should retain about 80 percent crown cover



Figure 5.—Shelterwood seed cut in northern hardwoods leaving approximately 70 percent crown cover (approximately 60 ft² of basal area per acre). Marking from below followed by brush saw removal of stems under 5.0 inches d.b.h. has created ideal conditions for regenerating tolerant and moderately tolerant species.

for hardwoods (60 to 70 ft²) and softwoods (100 to 120 ft²); for intermediately tolerant regeneration (chiefly yellow birch), the seed cut should leave a residual stand of about 30 to 50 percent crown cover (30 to 40 ft²); perhaps a little higher on wet sites. Marking for seed cuts (and preparatory cuts) must be from below, removing smaller stems as first priority, and leaving a uniformly distributed stand. Tables 17–19 in the Appendix help relate crown cover to basal area by species group. These tables allow shelterwood prescriptions to be written in terms of crown cover or basal area, or both.

The final removal cut for any species should be made when the regeneration is 3 to 4 feet tall or more for most species (> 2 feet for birch; > 1 foot for spruce). Winter removal minimizes logging damage to the regeneration. However, summer removal is a possibility with hardwoods

because of their sprouting ability. Other logging precautions to minimize damage to the regeneration include the careful layout of major skid trails, directional felling, log-length skidding, and the use of winching devices.

In previously cutover stands, where a good stocking of saplings and poles are present under an existing overstory, a natural shelterwood can be applied simply by removing the overstory. The main concerns are damage to, and adequate stocking in, the residual stand. Several planned modifications to the shelterwood system have been tried where the time between the seed cut and removal cut has been lengthened to maintain continual cover for esthetic purposes; these are known as delayed or extended shelterwoods. In the extreme, this approach becomes a two-age system where removal cuts are made at half-rotation intervals.

Dense, undesirable understory vegetation will hinder the establishment of regeneration under the shelterwood system. Methods for dealing with this problem include understory biomass operations, broadcast chemical treatments, and the mechanical or chemical treatment of individual unwanted stems.

The shelterwood system is a good option in mixedwood stands, especially those with high sawtimber potential. This system can be used to increase the tolerant softwood component; to maintain windfirmness if high residual crown cover is maintained; and to allow for the use of large equipment that tends to destroy understory saplings and poles.

Regeneration

Species favored by clearcutting and dense or open shelterwood cutting are summarized in Table 2. Clearcuttings commonly have 20,000 to 30,000 or more stems per acre 1 foot tall or taller at 5 years of age. Species composition is more important than numbers. About two-thirds of the milacres on clearcut areas generally are dominated with intolerant or intermediately tolerant species, though the proportion based on total numbers is less. If advanced regeneration of tolerants larger than 2 inches d.b.h. is retained, a somewhat lower proportion of intolerants-intermediates will result. Tolerant softwoods seldom regenerate well following clearcutting unless well-established regeneration is present—a type of natural or unplanned shelterwood.

One approach to evaluating regeneration following clearcutting is to take a series of circular plots, each 1/700 to 1/1,000 acre in size (8.9 to 7.4 feet in diameter). These plot sizes represent the area occupied by each tree when the stand reaches 4 to 6 inches d.b.h. (quadratic mean stand diameter or tree of average basal area) in the northern hardwood and mixedwood stocking guides (Figs. 6–7). Determine and record the dominant free-to-grow species—the species that will dominate the plot using all available information on tolerance, relative growth rate, longevity, etc. If the proportion of plots dominated by desirable species exceeds 60 percent (many plots contain more than one commercial stem), this would be equivalent to B-line stocking or better. Stocking of 40 percent would be about equivalent to C-line stocking. By also recording the desired species present that are not free to grow, it is possible to determine whether the species potential of each plot could be improved by a weeding operation. For example, a plot might be dominated by free-to-grow aspen; if removed, the plot might be dominated by paper birch. In summarizing the data, it is then possible to examine stocking of acceptable species both with and without treatment. If the without-treatment stocking is less than C line, but the with-treatment stocking is well above the C line, a weeding/cleaning operation should be considered.

Data from shelterwood cuttings in the Lake States indicate that at least 5,000 well-distributed seedlings per acre, 3 to 4 feet tall, should be present before the removal cut. After the removal cut, the regeneration can be evaluated using the plot system described for clearcutting.

In the past, scarification has been recommended for yellow and paper birches since most studies show much higher stocking of these birches on scarified seedbeds. However, scarification operations are expensive and difficult to justify. Recent experience indicates that summer logging, which encourages a small amount of scarification from the logging operation, does not necessarily produce more birch than winter logging.

Scarification during the seed cut of a shelterwood has been recommended in the Lake States for regenerating hemlock. In the Northeast, certain sites—notably the wetter ones—appear to develop a strong understory of hemlock without scarification. However, on drier mixedwood sites where little or no advance regeneration of hemlock is present, scarification during the shelterwood seed cut would appear to be helpful.

Planting is seldom done on a commercial scale in northern hardwoods and related types in the Northeast. However, where seed sources of desired species are lacking or genetically improved trees are desired, planting can be done successfully. We recommend container-grown stock for rapid growth and minimum mortality. The planting site needs to be freed of brush or sod—by mechanical or chemical means—to at least 3 to 4 feet around each seedling location. Posttreatment release often is necessary. In some areas, it also will be necessary to control damage from deer, rabbits, and mice. Yellow and paper birches, spruce, and hemlock all are possible planting candidates.

Stocking

Stocking guides for even-aged hardwood stands are given in Figure 6 and for mixedwood stands in Figure 7. The guides apply to the main crown canopy, i.e., excluding the suppressed trees. Mixedwood stocking applies to stands with 25 to 65 percent softwoods in the main crown canopy. The A lines represent the average density of undisturbed even-aged stands. The B lines represent the minimum density for maximum basal area or cubic-foot growth. The charts were developed from both simulated and remeasured plot data which show that maximum growth per acre occurs at about 55 to 65 ft² of basal area in hardwoods. The C line represents minimum stocking—the minimum amount of acceptable growing stock to make the stand worth managing. The C line is roughly 10 years' growth below the B line. Growth per acre (in basal area or cubic feet) is a little lower at the C line than the B line, and diameter growth more rapid.




Figure 6.—Stocking guide for main crown canopy of even-aged hardwood stands (beech-red maple, beech-birch-maple) shows basal area and number of trees per acre and quadratic mean stand diameter. The A line is fully stocked, the B line is suggested residual stocking. The C-line is minimum stocking. The quality line is the density required to produce high quality stems of beech, sugar maple, yellow birch, and red maple.

Average density of mixedwood stands is higher than that of hardwood stands (Figs. 6-7). The mixedwood A line is from 20 to 55 ft² above the hardwood A line. Similarly, the B line is from 35 to 45 ft² above the hardwood B line when the percentage of softwood is from 25 to 65 percent of the basal area of trees in the main crown canopy.

A Quality line also is shown for hardwoods (Fig. 6). Limited research indicates that species such as beech, sugar maple, red maple, and yellow birch do not prune well naturally unless grown at 80 ft² of basal area per acre. Paper birch, aspen, and white ash appear to be the only common hardwoods in the Northeast that will develop acceptable quality in small poletimber stands maintained at or near the

B line. At average stand diameters of about 6 inches, clear lengths of about 1 1/2 logs should be present on many trees. At this time or after additional clear bole development the stand can be thinned back to the B line (plus 5 to 10 ft² to allow for logging damage), perhaps in two operations if basal area is high and crowns are small. Up to stand diameters of roughly 6 inches, light improvement work to maintain species composition and select for stem quality is acceptable. The Quality line in Figure 6 is dotted because species composition and local experience will influence the level of stocking required in young stands to ensure quality development. To grow limb-free or small limbed hemlock and spruce, density should be maintained




Figure 7.—Stocking guide for main crown canopy of mixedwood stands (25 to 65 percent softwoods) shows basal area and number of trees per acre and quadratic mean stand diameter. The A line is fully stocked, the B line is suggested residual stocking, the C line is minimum stocking.

near the mixedwood A line for stands with a mean stand diameter less than 6 inches.

Growth, Yield, and Rotation

Board-foot and cubic-foot volumes per acre for unmanaged hardwood stands are given in Table 8, based on simulation (Solomon and Leak 1985). These are gross yields with no deduction for cull or defect. Cubic volumes of mixedwood are at least 15 to 25 percent greater than hardwood yields, and board-foot volumes may be proportionately even greater. However, precise estimates of mixedwood yields are not yet available. Yields of intensively managed hardwood stands are simulated in Table 9. These yields are for stands kept at 80 ft² or more of basal area until the mean stand d.b.h. reached 6 inches, and then thinned to B-line density when stand basal area was about two-thirds the

distance from B line to A line. Hardwood yields represent the maximum attainable under intensive silviculture in a natural stand and total managed yields at about 100 years of age are 50 to 90 percent greater than unmanaged yields. In applying both the managed and unmanaged yield tables, deductions from gross yields must be made for sound and rotten cull, logging waste, poor stocking, and nonforest acreage.

The site indexes in Tables 8 and 9 are for sugar maple, base age 50. Sugar maple sites of 70 and above commonly occur in New England on soils enriched by organic matter or derived from rich bedrock (e.g., limestone) or alluvium. Sugar maple and ash are common on such sites. Sites of 60 to 70 typically are beech-birch-maple sites; the soils are well- to moderately well-drained, fine-textured tills. Sites of

Table 8.—Volumes per acre in cubic feet (4.0-inch ib top) and board feet (8.0-inch ib top) for unmanaged hardwood stands, by site index, mean stand diameter, and age

Mean d.b.h. (inches)	Site 50		Site 60		Site 70				
	Age	Cubic feet	Board feet	Age	Cubic feet	Board feet	Age	Cubic feet	Board feet
	Years		Years		Years				
4.0	35		30		25				
6.0	59	1289	49	1547	41	1821			
8.0	83	1606	2983	67	1924	3560	55	2254	4258
10.0	120	1934	5554	87	2311	6640	69	2675	7632
12.0	182	2272	8259	114	2700	9783	85	3144	11461
14.0				157	3102	13048	102	3579	15079
16.0				196	3154	13257	127	3654	15390

50 to 60 tend to be beech-red maple or mixedwood sites; soil productivity is limited somewhat by shallow, compacted layers, coarse textures, restricted aeration, or excessive stoniness. Sites below 50 usually are poorly drained or shallow to bedrock. Sites below about 50 to 55 are best suited to growing softwoods, or hardwoods on shorter rotations (e.g., paper birch and aspen).

Rotation age usually is based on (1) the culmination of mean annual increment, or (2) the time required to grow a certain-size tree or product. Tables 8 and 9 indicate that culmination of mean annual board-foot growth in both managed and unmanaged stands ranges from about 100 to 120 years. Mean diameters at the point of culmination vary with site—the range is 14 to 18 inches. Culmination of mean annual increment for cubic volume occurs at age 40 to 50 in unmanaged stands and 80 to 90 years in managed stands. In Table 9, some inconsistency is evident in the trend of mean annual increment because of the timing of intermediate cuts.

Intermediate Cuttings

Where there are good markets for pole-size material, non-commercial thinning/cleaning often is not needed; in many cases, silvicultural needs can be met through commercial operations in pole stands. However, if an analysis of the reproduction following clearcutting or shelterwood cutting indicates that desired species objectives will not be met (see section on even-age regeneration), a noncommercial operation may be warranted. The silvicultural objective should be to increase the proportion of plots dominated by acceptable species to about 40 percent (equivalent to C-line stocking). Where the objective is to increase the proportion of softwood species, cleaning can be done in seedling stands with selective herbicides. To change the species mix in hardwood stands, mechanical or chemical

stem treatments should be done in stands between about 10 to 20 years of age. Examples where noncommercial work might be warranted are in: (1) mixedwood regeneration where the objective is to grow softwoods; (2) mixtures of valuable hardwoods (yellow birch, sugar maple, ash) in combination with fast-growing less valuable species such as red maple; (3) other situations where economic analysis indicated that costs are justified.

In most hardwood stands between 4 and about 6 inches mean d.b.h., the stocking guide (Fig. 6) recommends fairly high stocking for those species that are resistant to natural pruning. Improvement work during this period might be accomplished by light, commercially marginal operations that remove 15 to 25 ft² of basal area per acre. In stands of paper birch, ash, and aspen averaging 4 to 6 inches d.b.h., heavier cuttings down to the B line are permitted for fuelwood or pulp.

Pruning is not a common silvicultural treatment in northern hardwoods. But the high value differential between clear and knotty logs is reason enough to continue to exam the prospective costs and returns from this practice. Pruning probably is most feasible for valuable species that are moderate to poor self-pruners: sugar maple, yellow birch, and red oak. Prune trees that are about 4 to 6 inches during the late summer or dormant season; do not remove more than a third of the live crown. Do not flush cut. Place the saw just outside the branch bark ridge and cut downward and slightly outward.

Table 9. Volumes per acre in cubic feet (4.0-inch top) and board feet (8.0-inch top) for intensively managed even-aged hardwoods

Mean d.b.h. (inches)	Site 50						Site 60						Site 70					
	Cumulative thinnings		Standing		Total		Cumulative thinnings		Standing		Total		Cumulative thinnings		Standing		Total	
	Cubic feet	Board feet	Cubic feet	Board feet	Cubic feet	Board feet	Cubic feet	Board feet	Cubic feet	Board feet	Cubic feet	Board feet	Cubic feet	Board feet	Cubic feet	Board feet	Cubic feet	Board feet
4.0																		
6.0	226	1158	2008	2698	1384	2008	269	1418	1189	2211	2432	3106	315	1657	1426	2760	2865	3835
8.0	1012	996	2620	5420	2620	5420	1243	1912	1854	3893	10005	6366	1439	1075	2222	6400	3661	7475
10.0	1012	1608	4694	8620	3245	8620	1854	2039	7375	10005	68	2148	3171	2395	8807	4543	11978	
12.0	1520	1725	6409	12098	3866	12098	2602	2011	8449	14082	77	3019	6650	2389	10139	5408	16789	
14.0	2128	4596	1738	7502	6340	4255	2602	5633	10289	5051	15922	85	3019	6650	2868	12203	5887	18853
16.0	2786	7435	1469	6340	4255	13775	3394	8960	8760	5479	17720	92	3915	10465	2446	10446	6361	20911
18.0																		

Years
35
57
74
89
105
124
147

Years
30
48
61
72
83
95
107
119

Years
25
40
50
59
68
77
85
92

Beyond 6 inches mean diameter, commercial thinnings generally will be feasible, lowering the basal area to about the B-line level (perhaps in two operations) plus an allowance of 5 to 10 ft² for logging damage. Then, when a stand reaches one-half to three-fourths of the distance from the B line to the A line, additional commercial thinnings can be made to reduce the basal area back to B-line level (plus damage allowance). Most commercial thinning will be in the main crown canopy, removing dominant, codominant, and intermediate trees. Keep in mind that certain types of marking (from above, from below) may change the residual mean diameter and also the appropriate B line. The objective is to provide adequate growing space for the stems with highest value potential by removing:

1. Risk trees: Valuable trees that will not last until the next thinning, or that will experience severe degrade.
2. Unacceptable stems: Trees that will not produce sawlog material now or in the future due to defect or cull.
3. Undesirable species.
4. Acceptable stems crowding high-value stems.

Stand Evaluation

Reproduction and Sapling Stands

In these young even-aged stands (mean stand diameter up to 4.0 inches) the primary need is for a method of judging the adequacy of stocking and species, and predicting the need for early noncommercial treatment.

To determine stocking, sample about 2 plots per acre in each young stand up to a total of about 50; plot size should be 8.9 feet (1/700 acre) or 7.4 feet (1/1,000 acre) in diameter. Record:

1. The species that will dominate the plot if left untreated. This requires the application of all available knowledge on species growth rates, tolerance, longevity, etc.
2. The desirable species not free to grow (commercial or desirable species) that will dominate the plot if one or two undesirable overstory stems are removed.

If at least 40 to 60 percent of the plots are dominated by desirable free-to-grow stems, the stand should attain C-line or B-line stocking of acceptable species when it reaches the lower end of the stocking guide. If stocking of desirable free-to-grow stems is less than 40 percent, the stocking of

desirable species not free to grow should be examined to determine whether a precommercial operation will raise the representation of desirable species to C-line or B-line levels.

Poletimber and Sawtimber Stands

These are even-aged or uneven-aged stands with mean diameters larger than 4.0 for trees in the main crown canopy. Take a minimum of 10 systematically located sample points in uniform stands, and up to 30 points in variable stands. On a cumulative tally (Table 10) (or a conventional tally and with the data in Table 11) record trees counted with a 10-factor prism by 2-inch diameter classes, and the following tree classes (denoted by the tally legend):

1. Species or species group (optional)
2. Acceptable growing stock
 - a. Mature trees (optional if species are tallied)
 - b. Immature trees (optional)
3. Unacceptable stems
 - a. Defective (optional)
 - b. High risk (optional)
 - c. Cull (optional)

For uneven-age management, the tally should include all trees in the 6-inch class and larger. For even-age management, the tally should include all trees in or touching the main crown canopy (exclude the suppressed trees). Where the choice has not yet been made between even-age and uneven-age management, the tally legend should distinguish between suppressed trees and those in the main crown canopy. Acceptable growing stock will produce sawlog or better material now or in the future. Unacceptable stems will not. Maturity can be tallied in the field using the size guidelines in Table 5, and current tree condition can be noted. If the tally legend separates species or species groups, maturity can be scored later using the general guidelines at the bottom of Table 12. Also, in even-age stands, measure breast height age and total height for up to five dominant stems per stand to determine site index (Figs. 8-9). Determine whether the stand is beech-birch-maple, beech-red maple, or mixedwood. And judge on the ground whether a commercial cutting is now feasible; this judgment should be based on volume, quality, accessibility, and markets.

Table 10. Sample cumulative tally for a 10- or 20-factor prism (example for two plots)

No. trees	Diameter at breast height (inches)										Tally	No. Plots		
	10	20	2	4	6	8	10	12	14	16			18	20
1	458	115	51 ^X	29 ^o	18 ^o	13 ^o	9 ^o	7 ^o	6 ^M	5	4	3	3	3
2	917	229	102 ^X	57 ^o	37 ^o	25 ^o	19 ^o	14 ^o	11 ^o	9	8	6	5	5
3	1375	344	153	86 ^o	55 ^o	38 ^o	28 ^o	21 ^o	17	14	11	10	8	8
4	1834	458	204	115	73 ^o	51 ^o	37 ^o	29	23	18	15	13	11	11
5	2292	573	255	143	92 ^o	64	47	36	28	23	19	16	14	14
6	2750	688	306	172	110	76	56	43	34	27	23	19	16	16
7	3209	802	357	201	128	89	65	50	40	32	27	22	19	19
8	3667	917	407	229	147	102	75	57	45	37	30	25	22	22
9	4125	1031	458	258	165	115	84	64	51	41	34	29	24	24
10	4584	1146	509	287	183	127	94	72	57	41	34	29	24	24
11	5042	1260	560	315	202	140	103	79	62	47	37	30	25	25
12	5501	1375	611	344	220	153	112	86	68	51	41	34	29	29
13	5959	1490	662	372	238	165	122	93	74	57	45	37	30	30
14	6417	1604	713	401	257	178	131	100	79	62	51	41	34	34
15	6875	1719	764	430	275	191	140	107	85	67	53	43	35	35
Trees/acre:			51	43	46	25.5	18.5	10.5	5.5	5.5	5.5	5.5	5.5	5.5
BA/acre:			10	15	25	20	20	15	10	10	10	10	10	10

Trees/acre: Add the last numbers used in each column and divide by the number of plots.

BA/acre: Add the number of entries; multiply by the BA factor; and divide by the number of plots.

BA factor 10: Use the numbers consecutively in each column. For example, one 10-inch tree in a plot represents 18 trees per acre; two 10-inch trees represents 37 trees.

BA factor 20: Use the second number of each pair in a column. For example, one 10-inch tree in a plot represents 37 trees per acre; two 10-inch trees represents 73 trees.

Site-Index Trees:

Species	Age	Height	Type
SM	70	70	Beech-birch-maple

Total = 149
Total = 105

No. Plots

Legend
/ Acceptable
0 Unacceptable
X Suppressed
M Mature

Table 11.—Basal area per tree and numbers of trees per acre conversion for a 10-factor prism

D.b.h.	Prism conversion	Basal area per tree	D.b.h.	Prism conversion	Basal area per tree	D.b.h.	Prism conversion	Basal area per tree
<i>Inches</i>	<i>No. trees/acre</i>	<i>Ft²</i>	<i>Inches</i>	<i>No. trees/acre</i>	<i>Ft²</i>	<i>Inches</i>	<i>No. trees/acre</i>	<i>Ft²</i>
1.0		0.0055	11.5		0.7213	22.0	3.8	2.6398
1.5		.0123	12.0	12.7	.7854	22.5		2.7612
2.0	458.4	.0218	12.5		.8522	23.0	3.5	2.8852
2.5		.0341	13.0	10.8	.9218	23.5		3.0121
3.0	203.7	.0491	13.5		.9940	24.0	3.2	3.1416
3.5		.0668	14.0	9.4	1.0690			
4.0	114.6	.0873	14.5		1.1467			
4.5		.1104	15.0	8.2	1.2272			
5.0	73.3	.1364	15.5		1.3104			
5.5		.1650	16.0	7.2	1.3693			
6.0	50.9	.1963	16.5		1.4849			
6.5		.2304	17.0	6.3	1.5763			
7.0	37.4	.2673	17.5		1.6703			
7.5		.3068	18.0	5.7	1.7671			
8.0	28.6	.3491	18.5		1.8667			
8.5		.3941	19.0	5.1	1.9689			
9.0	22.6	.4418	19.5		2.0739			
9.5		.4922	20.0	4.6	2.1817			
10.0	18.3	.5454	20.5		2.2921			
10.5		.6013	21.0	4.2	2.4053			
11.0	15.2	.6600	21.5		2.5212			

The essential information from the plots can be summarized (Table 12) to provide a basis for either the uneven-age or even-age stand options. For the uneven-age summary, the first six basal area columns provide a description of the initial stand; not all columns need be used, or more can be added to provide a species breakdown. From these data, the initial percentage of sawtimber can be determined, as can the initial approximate q from the tabulation in the section on uneven-age stocking. If the prescription key suggests a harvest cutting, the residual goal is determined by: (1) examining various approaches (marking rules) for removing the poorer quality material so as to leave a good-quality stand with the required total basal area; or (2) using a residual goal based on the initial q of the stand; or (3) using the general guidelines in Table 6. The marking goal is simply the difference between the initial total basal area and the residual goal; however, 5 to 10 ft^2 may be subtracted from the marking goal for logging damage.

For the even-age summary, basal area of the initial stand is listed by tree condition class. Number of trees per acre is taken from the cumulative tally. Total basal area per acre

and number of trees are used to read mean stand diameter from the stocking chart (Figs. 6–7). Basal area at the A, B, C, and Quality lines also are taken from the stocking chart. If the prescription key calls for a treatment, the residual goal generally is determined by the B line or Quality line. However, residual goals higher than the B line may be prescribed to maintain maximum amounts of quality material, for esthetic purposes, etc. The marking goal is the difference between the initial and residual, minus any allowance for logging damage. Distributing the residual goal and marking goal among tree condition classes helps in the development of marking guides and helps ensure that the treatment will improve the quality of the stand.

Stand Prescription

Key

Use the following key to identify the stand condition and find the appropriate prescription (A, B, C, etc.). Details of the prescriptions follow the key. Also, consult the appropriate section describing the treatment within the text.

Table 12.—Summary table for uneven-aged or even-aged stand diagnosis (example from Table 10)

D.b.h. class (inches)			Defective	High risk		Cull	Total	Possible residual goals (Q = 1.5-1.7)		Possible marking goals	
	Mature	Immature		risk	Cull			Q = 1.5	Q = 1.7	goals	goals
-----Ft-----											
UNEVEN-AGED											
6-10		40	10				50	30	38	20	12
12-14		10	30				40	20	18	20	22
16+	5	5	15				25	20	14	5	11
All	5	55	55				115	70	70	45	45
Initial % Sawtimber = 57						Initial Q = 1.5					
-----Ft-----											
EVEN-AGED											
-----Ft-----											
Tree condition		Initial stand		Residual goal		Marking goal					
-----Ft-----											
Mature		5		5		0					
Immature		45		45		0					
Defective-high risk		55		14		41					
All		105		64		41					
No. Trees		149		Commercial Cutting		Mature D.b.h. (inches)					
MSD		11.4		___ Feasible		20 - SM, YB, WP, Hem					
A line BA		112		___ Not feasible		18 - Be, WA, RO					
B line BA		64								16 - RS, RM	
C line BA		48								12 - PB, Asp., BF	
Quality line BA		___									
Bole Condition:		___ Clear merchantable length		Site-Index Trees							
		___ More natural pruning needed		Species						B ___	
				Age						70 ___	
				Height						70 ___	
				Site						60 ___	

Reproduction or Sapling Stands (Mean d.b.h. of Overstory Less Than 4.0 Inches)

1. 40 percent or more of the plots stocked with a desirable free to grow stem (untreated). A

1. Less than 40 percent of the plots stocked with a desirable free to grow stem (untreated).

2. More than 40 percent, preferably more than 60 percent, of the plots stocked with a desirable stem not free to grow. B

2. Less than 40 percent of the plots stocked with a desirable stem not free to grow. C




Figure 8.—Site-index curves (breast height age 50) for paper birch, white ash, yellow birch, and sugar maple in Vermont and New Hampshire. Values in parentheses are for site-index breast height age 75 (Curtis and Post 1962b).

Poletimber and Sawtimber Stands (mean d.b.h of overstory 4.0 inches or more)

1. Objective: uneven-age management
2. Acceptable mature and immature growing stock more than:
 - 40 ft² (hardwood stand) or
 - 60 ft² (mixed-wood stand)
3. Acceptable mature and immature growing stock 12 inches and larger more than:
 - 25 ft² (hardwood) or
 - 40 ft² (mixedwood)
4. Total basal area more than:
 - 100 ft² (hardwood) or
 - 130 ft² (mixedwood)
4. Total basal area less than:
 - 100 or 130 ft²

D
E

3. Acceptable mature and immature growing stock 12 inches d.b.h and larger less than:
 - 25 ft² (hardwoods) or
 - 40 ft² (mixedwood)
4. Total basal area more than:
 - 100 ft² (hardwood) or
 - 130 ft² (mixedwood)
4. Total basal area less than:
 - 100 or 130 ft²
2. Acceptable mature and immature growing stock less than:
 - 40 ft² (hardwood) or
 - 60 ft² (mixedwood)
1. Objective: even-age management
2. Stocking of acceptable growing stock less than C line for the appropriate type

F
G
H
I




Figure 9.—Relationships among site indices (base age 50) for white ash, paper birch, yellow birch, and sugar maple. To estimate site index of species X from site index of species Y: find known site index on curve for species Y; move vertically up or down to curve until species X is located; read horizontally across to the left to find estimated site index for species X (Curtis and Post 1962a).

2. Stocking of acceptable growing stock more than C line
3. Stand mature^{1/}
 4. Objective: intolerant and intermediate species J
 4. Objective: tolerant species K
3. Stand not mature
 4. Stand more than 6 inches mean stand diameter and adequate clear length developed
 5. Total stocking more than halfway between A and B lines

¹Mature: (1) at rotation age, (2) at mature size, based on product objectives, or (3) 50 percent or more of the basal area in mature trees.

6. Commercial thinning feasible now or within 10 years L
6. Commercial thinning not feasible now or within 10 years M
5. Total stocking less than halfway between A and B lines N
4. Stand less than 6 inches mean stand diameter, or clear length not well developed
5. Total stocking more than Quality line
6. Light commercial thinning feasible now or within 10 years O
6. Light commercial thinning not feasible now or within 10 years P
5. Total stocking less than Quality line Q

Prescriptions

- A. This young stand should develop naturally into an adequate pole stand with at least C-line stocking of desirable species. Noncommercial treatment is not required.
- B. This young stand will develop into a pole stand with C line stocking of desirable species only if cleaned precommercially. Benefits and costs of such treatment should be examined using all available information on site, species response, and management objectives. Apply the treatment if warranted.
- C. Even if cleaned precommercially, this young stand will not develop C-line stocking of desirable species. Reexamine the stand in 10 to 20 years to determine the best treatment options—including the possibility of biomass harvesting for fuel or fiber.
- D. This stand has suitable quality, structure, and basal area to implement uneven-age management. Develop and apply marking guides to meet goals for residual basal-area structure, tree condition, and regeneration. Consider both single-tree and group selection.
- E. This stand has suitable quality and structure to implement uneven-age management. But stand density is not critically high. Reexamine in 10 to 20 years, unless the possible loss of valuable high-risk trees warrants immediate harvest cut by selection or group-selection methods.
- F. This stand has suitable quality and density to initiate uneven-age management, but sawtimber stocking is low. Apply a commercial improvement cut, removing lower quality overstory stems, leaving a residual basal area of about 65 to 70 ft² (hardwood) or 80 to 100 ft² (mixed wood) per acre plus any allowance for logging damage.

- G. This stand has suitable quality for uneven-age management, but sawtimber stocking is low and stand density is not critically high. Reexamine in 10 to 20 years.
- H. This stand has too little quality growing stock for efficient uneven-age management. Reconsider the possibility of even-age management through clearcutting or shelterwood cutting. The other alternative is a long series of improvement cuts and selection/group selection to gradually improve the condition of the stand.
- I. Acceptable growing stock is inadequate. Plan to regenerate the stand with clearcutting, strip cutting, or shelterwood cutting when commercially feasible.
- J. Apply clearcutting to maximize the proportion of intolerant and intermediate species. Strip cutting should maximize intermediates such as yellow birch. In sensitive areas, a heavy two-cut shelterwood can be applied by leaving 30 to 50 percent residual crown cover (30 to 40 ft²) following the seed cutting and removing the overstory in about 5 years.
- K. Use a light two-cut shelterwood, leaving about 80 percent or more crown cover (60 to 70 ft² of basal area), during the initial seed cutting and removing the overstory when the tolerant advanced regeneration is more than 3 feet tall.
- L. This immature stand has adequate young growing stock for even-age management, and sufficient stand density to support a commercial thinning. Stands should be thinned to not below the B line. However, only up to one third of the main canopy basal area should be removed at any one time. In stands within about 20 years of maturity, commercially thin only if there will be losses in volume or value if the stand is left untreated until final harvest.
- M. This immature stand has adequate acceptable growing stock and density for even-age management, but commercial thinning is judged not feasible because of accessibility, current markets, etc. Leave untreated until commercial thinning prospects improve.
- N. This immature stand has adequate acceptable growing stock for even-age management, but stand density is not critically high. Reexamine in 10 to 20 years.
- O. This immature stand has sufficient potential quality and density for even-age management, but adequate clear length has not yet developed. Light thinning or improvement cutting to the Quality line, removing a small amount of poor quality or risky material, is permitted; this option is best suited to stands where quality, species, and site index are above average.

- P. This immature stand has sufficient quality and density for even-age management, but adequate clear length has not yet developed. Light thinning is judged not feasible. Leave untreated, and reexamine in 10 years.
- Q. This immature stand has sufficient potential quality for even-age management, but adequate clear merchantable length has not yet developed. For production of quality material, leave the stand untreated so that increasing stand density will encourage natural pruning. For fuelwood production, the stand may be thinned to B line.

Regulation

Regulation refers to the methods used to control the amount and periodicity of timber yields from a property. Commercial timberland owners, industrial owners, and certain large public ownerships may need regular, sustained or increasing yields. Owners of small tracts may have less need to control yields.

With uneven-age management, periodic yields from each stand or group of stands are achieved by setting a residual stand density, structure, and growing-stock condition (in terms of risk and quality potential) that will produce good volume or value growth over the cutting cycle (see Tables 3 and 4).

The first cut in a heavily stocked stand will produce fairly high gross yields, but may be low in net yield and value. Ensuring harvests in any stand are made at intervals equal in length to the cutting cycle. During these harvests, residual stand density is roughly consistent, though the proportion and quality of the residual sawtimber may be increased gradually until it reaches a desirable level. This approach will result (after the first cut) in fairly constant cubic-foot yields roughly equal to annual growth times the cutting cycle, and gradually increasing sawtimber yields until an essentially constant level is reached.

On a large uneven-aged property, where annual yields are feasible and desired, the entire property can be divided into a number of cutting units or groups of stands equal to the years in the cutting cycle. Then, each year, a different cutting unit is harvested to provide an annual yield. At the outset, units are entered in order of priority based on maturity, risk, stocking, etc.

Uneven-age regulation commonly is called volume, basal area, or growing-stock control. However, since the cutting units will have roughly equal acreages (or acreages inversely proportional to productivity), there is some element of area control involved as well.

The system becomes more complicated when there are inaccessible or less productive areas on a long cutting cycle, and productive or accessible areas on a short cutting cycle. Detailed scheduling is required to assure that roughly equal yields are harvested each year. A more formal approach is to divide the inaccessible lands into a number of cutting units equal to the number of years in the long cutting cycle, and also to divide the accessible lands into a number of cutting units equal to the years in the short cutting cycle. Then, each year, both an inaccessible and an accessible unit are cut. This approach tends to regulate both yields and access costs.

With even-age management, there are two components to the yield: harvest-cutting and thinning yields. In theory, an even-aged forest is fully regulated when it has roughly equal acreages in each 10- or 20-year stand age class from the youngest class up to the class representing the planned rotation age. As with uneven-age management, it sometimes is practical to: (1) divide the entire property into type or accessibility classes, (2) set an appropriate rotation age and thinning interval for each class, and (3) work toward an balanced age distribution in each type of accessibility class. To develop a balanced age distribution, harvest an acreage per year equal to the total acreage divided by the planned rotation age. If the entry period for harvest cutting is more than 1 year, multiply by the number of years in the entry cycle to determine the acreage to harvest at each entry.

Thinned acreage commonly is 2 or 3 times the harvest acreage, though thinned volume may be between 50 and 100 percent of the harvest yields. In an unbalanced even-aged forest, the thinning yields will be variable because: (1) the acreage in each age class will be unequal, and (2) the quality and stocking of some acres in each age class may not warrant thinning.

Economic Considerations

With moderately intensive silviculture, managed stands can yield at least 50 percent more volume than unmanaged stands. Although the increased physical yields seem worth pursuing, the financial returns from those yields may not be. The following discussion explores the economic effectiveness of applying silvicultural guidelines in the management of northern hardwood forests.

We develop a generalized case to trace the changes in timber values that we might expect in northern hardwood stands over long periods of time under various management strategies. We also assign estimates of stumpage values and their costs to the volume yields indicated in Appendix Tables 20-23, and then compare the resulting timber values. Methods for testing the economic effective-

ness of managing a particular stand are available elsewhere (Leak 1980).

Hardwood Diversity

In any discussion of northern hardwoods and associated types, we must first emphasize their diversity. Each species has its own package of characteristics as to strength, workability, appearance, and appeal. Market prices attest to this and tend to differentiate relative values among species.

Eastern hardwood stands also are diverse in product potential. Figure 10 depicts a hierarchy of relative product values, along with a general woods-run volume distribution that is typical of many eastern hardwood stands. Although the actual values vary from one stand to another because of species and size mix, logging conditions, and markets, the relative value differences among products often are large (DeBald 1981).

Timber size and quality are especially important in northern hardwoods. Figure 11 shows the relative values of trees by both diameter and butt-log grade. The increased value through increased size suggests concentrating growth on selected fast-growing crop trees. The large differences in tree values from one butt-log grade to another indicate the importance of concentrating growth on trees that are likely to improve in grade.




Figure 10.—Typical value/volume hierarchy, eastern hardwood stands.




Figure 11.—Relative values of two-log hard maple trees, by butt-log grade.

Potential Timber Values

The biological recommendations in this silvicultural guide capitalize on the diversities mentioned through the manipulation of species mix, tree size, timber quality, and product objectives. But are higher timber values worth waiting for? Are they worth working for? The short answer is: probably. The long answer depends on a number of factors.

One factor concerns the anticipated timber yields themselves. As shown earlier, we can expect timber yields sooner through silvicultural activities. We also can expect the overall volume yields to be greater. But in order to test the economic effectiveness of our silvicultural guidelines, we need to assess the possible dollar returns from those physical yields, along with their costs.

We must first consider the product potential of a stand with and without silvicultural treatment (Table 13). We would, for example, expect a low product potential in unmanaged stands and higher potentials in managed stands, depending on our efforts to develop those potentials. In Table 13, product distribution A represents a typical unmanaged

northern hardwood stand (Filip and Williams 1968). Distributions B, C, and D represent a range of improved product mixes that might be expected through the application of silvicultural guidelines, and reflect an upgrading of timber quality through thinnings.

Table 13.—Assumed percentages of sawtimber volume

Product	Product distribution			
	A	B	C	D
Veneer	2	4	6	8
Sawlogs				
High quality	3	6	9	12
Medium quality	40	45	50	55
Low quality	15	15	15	15
Pallet stock	40	30	20	10

Next, we must assign dollar values to potential products. Recent stumpage prices in New Hampshire indicate a wide range in values by product class and highlight the value premiums for higher value products (Table 14). They also point to the importance of encouraging the development of high-value species.

To evaluate the development of quality over long and varying timespans, we should include price-change expectations in our valuations. Let's use real rates of price change—rates over and above inflation. Using real prices, eliminates the need to make an additional guess at expected long-run inflation rates and reflects the fact that hardwood lumber-price trends often exceed inflation rates. Let's use the stumpage prices in Table 14 for a base and project them to increase in real value as follows:

Log quality	Real rate of increase (%)
Veneer and high	3
Medium	2
Low	1
Pallet stock	0.5

We can then assign estimates of dollar values to both the thinnings and the standing volumes shown in Tables 20-23—in real terms. The results will portray the benefits of quality development and our expectations of how silvicultural treatments can enhance the development of quality.

But every benefit has its cost. We need, then, to also consider the costs of maintaining and managing a timber stand—again, over long and varying timespans. So, let's assume an annual real cost of \$1 per acre for such things as property taxes and maintaining boundaries. Although the annual cost is common to each of the management

strategies, the total costs will vary depending on the timespans that the management strategies cover. Let's assume, further, that consulting forester fees account for 10 percent of timber-sale proceeds, a 30-percent income tax bracket, and capital gains treatment of 40 percent.

Net Present Values

But we also must consider the time value of money. For most of us, a dollar in hand today is worth more than a dollar to be received (or spent) 5 years from now. How much more depends on the rate of return (cost of capital) that we assume. The values in Table 15, for example, are net present values, the result of discounting expected net future values and expected future costs all back to year zero at a rate of 4 percent, then subtracting the discounted costs from the discounted values. Discounting to year zero converts varying timeframes to one common point in time and allows us to analyze stand-value development over time. The resulting net present values express all amounts in equivalent terms—today's dollars at time zero.

Comparison of different thinning regimes suggest that silvicultural activities aimed at improving product mix can result in a substantial increase in value yields (Table 15). The more we improve the potential product mix, the greater the value yield.

Although much of the increased value results from improved product mix, the timing of value yields also is important. The time required to reach a given mean stand diameter is considerably shorter with management than it is without management. Holding costs, then, are lower. But more important, the discounting period for managed stands of a given mean diameter are much shorter than they are for unmanaged stands of the same diameter. Note that even if product mix were not improved by thinnings (Table 15, Column A), the net present values of managed stands are much higher than those for unmanaged stands. The time value of money is extremely important.

The value of timber removed in thinnings also is important. The cash flows that they generate add greatly to the overall value yields from managed stands. In many managed stands, the net present value of timber sold from thinnings amounts to almost as much as the net present value of the standing timber.

The prospect of building up higher timber values faster, combined with cash flows from thinnings, suggests that managing northern hardwoods can be worthwhile. Depending on the degree of improvement in product mix, the value yields of managed stands can be dramatically higher than those of unmanaged stands.

Table 14.—Recent typical sawtimber stumpage prices (adapted from Engalichev 1984)

Product class	White ash	Hard maple	Yellow birch	White birch	Red oak	Other
	—Dollars/M bf—					
Veneer	135	90	115	110	170	45
Sawtimber						
High quality	120	90	105	85	135	40
Medium quality	90	65	80	65	100	30
Low quality	70	50	65	55	75	25
Pallet stock	25	25	25	25	25	25

Table 15.—Estimates of net present value for northern hardwoods by thinning regime^a and product distribution (based on projected real stumpage prices and 4-percent discount rate)

Mean d.b.h. (inches)	Stand age	Product distribution ^b			
		A	B	C	D
Years -----Dollars-----					
9-Inch Thinning					
8	67	5	12	19	25
10	83	20	33	46	58
12	98	35	54	74	93
14	110	42	66	89	113
16	125	35	58	81	104
18	142	32	55	78	101
7-Inch Thinning					
8	64	14	23	32	41
10	76	36	52	68	85
12	90	50	73	95	118
14	101	66	96	125	155
16	114	60	89	118	147
18	128	57	86	116	145
Quality-Line Thinning					
8	61	20	30	40	50
10	72	48	67	86	105
12	83	66	92	117	143
14	95	77	108	139	170
16	107	75	108	140	173
18	119	69	101	133	165
Unmanaged					
8	67	5	—	—	—
10	87	11	—	—	—
12	114	6	—	—	—
14	157	-6	—	—	—
16	196	-15	—	—	—
18	230	-19	—	—	—

^a Thinnings beginning at 9, 7, and approximately 5 (Quality line) inches mean stand diameter, and no thinning (unmanaged), and with yield schedules as shown in Tables 20-23.

^b See Table 13 for product distributions.

Rate of Return

As an alternative to net present value, we might consider a rate of return analysis of timber management strategies. The internal rate of return (IRR), for example, is the compound rate of interest that equates the present value of expected future returns with the present value of expected

future costs. It is the interest rate at which net present value is zero.

Using the same timber value and cost information that we used to estimate net present values, we estimated the internal rates of return for the same management strategies and product distributions. We found that we might expect managed northern hardwood stands to yield real rates of return that range from 5 to 8 percent (Table 16); and that unmanaged northern hardwoods might, at best, yield rates below 5 percent.

Note that the IRR cited are real rates. They do not include the effects of inflation. We can, though, approximate nominal or market rates by adding our inflation expectations to

Table 16.—Estimates of real rate of return for northern hardwoods by thinning regime^a and product distribution (based on real stumpage prices)

Mean d.b.h. (inches)	Stand age	Product distribution		
		B	C	D
Years -----Percent-----				
9-Inch Thinning				
8	67	5.0	5.3	5.6
10	83	5.5	5.8	6.1
12	98	5.7	5.9	6.2
14	110	5.7	6.0	6.2
16	125	5.6	5.8	6.0
18	142	5.4	5.7	5.9
7-Inch Thinning				
8	64	5.7	6.2	6.5
10	76	6.4	7.0	7.3
12	90	6.5	7.0	7.3
14	101	6.5	6.8	7.0
16	114	6.3	6.8	7.0
18	128	6.2	6.5	6.7
Quality-line Thinning				
8	61	6.2	7.1	7.5
10	72	6.8	7.2	7.5
12	83	6.9	7.5	7.8
14	95	6.9	7.4	7.7
16	107	6.7	7.0	7.3
18	119	6.6	7.0	7.2

^a Thinnings beginning at 9, 7, and approximately 5 (Quality line) inches mean stand diameter, and with yield schedules as shown in Tables 20-23.

the real rates.² For example, if we expected a 7 percent rate of return over over a span of years, along with an average inflation rate of 4 percent, the nominal IRR would be approximately 11 percent. The tough (if not impossible) part of making the conversion is trying to predict future inflation rates.

The differences in possible value yields, with and without management, seem wide enough to demonstrate the economic effectiveness of adopting silvicultural guidelines in the management of northern hardwood forests generally, and to warrant taking a closer look at the management potentials of individual stands, specifically.

Literature Cited

- Curtis, Robert O.; Post, Boyd W. 1962a. **Comparative site indices for northern hardwoods in the Green Mountains of Vermont**. Stn. Pap. NE-171. Upper Darby, PA: U.S. Department of Agriculture, Forest Service, Northeastern Forest Experiment Station. 6 p.
- Curtis, Robert O.; Post, Boyd W. 1962b. **Site index curves for even-aged northern hardwoods in the Green Mountains of Vermont**. Bull. 629. Burlington, VT: University of Vermont Agricultural Experiment Station. 11 p.
- DeBald, Paul S. 1981. **Estimating timber stand values: A guide to treatment**. In: Proceedings of the national silviculture workshop; Roanoke, VA; June 1-5, 1981; Washington, DC: U.S. Department of Agriculture, Forest Service. p. 237-247.
- Engalichev, Nicolas. 1984. **New Hampshire forest market report, 1983**. Ext. Publ. 11. Durham, NH: University of New Hampshire, Cooperative Extension Service. 34 p.
- Filip, Stanley M.; Williams, Frank W. 1968 **Volume and value of products harvested from a northern hardwood clear cutting**. Northern Logger. 17(4): 22-23.
- Gilbert, Adrian M.; Jensen, Victor S. 1958. **A management guide for northern hardwoods in New England**. Stn. Pap. NE-112. Upper Darby, PA: U.S. Department of Agriculture, Forest Service, Northeastern Forest Experiment Station. 22 p.
- Kingsley, Neal P.; Birch, Thomas W. 1977. **The forest-land owners of New Hampshire and Vermont**. Resour. Bull. NE-51. Upper Darby, PA: U.S. Department of Agriculture, Forest Service, Northeastern Forest Experiment Station. 47 p.
- Leak, William B. 1980. **Rapid economic analysis of northern hardwood stand improvement options**. Res. Note NE-296. Broomall, PA: U.S. Department of Agriculture, Forest Service, Northeastern Forest Experiment Station. 6 p.
- Leak, William B.; Solomon, Dale S.; Filip, Stanley M. 1969. **A silvicultural guide for northern hardwoods in the Northeast**. Res. Pap. NE-143. Upper Darby, PA: U.S. Department of Agriculture, Forest Service, Northeastern Experiment Station. 34 p.
- Shigo, Alex L. 1983. **Tree defects: a photo guide**. Tech. Rep. NE-82. Broomall, PA: U.S. Department of Agriculture, Forest Service, Northeastern Experiment Station. 167 p.
- Shigo, Alex L.; Larson, Edwin H. 1969. **A photo guide to the patterns of discoloration and decay in living northern hardwood trees**. Res. Pap. NE-127. Upper Darby, PA: U.S. Department of Agriculture, Forest Service, Northeastern Forest Experiment Station. 100 p.
- Solomon, Dale S. 1977. **The influence of stand density and structure on growth of northern hardwoods in New England**. Res. Pap. NE-143. Upper Darby, PA: U.S. Department of Agriculture, Forest Service, Northeastern Experiment Station, 34 p.
- Solomon, Dale S.; Leak, William B. 1985. **Simulated yields for managed northern hardwood stands in New England**. Res. Pap. NE-578. Broomall, PA: U.S. Department of Agriculture, Forest Service, Northeastern Forest Experiment Station. 24 p.

²Actually, we would use the formula $(1+n)=(1+r)^i(1+i)$ when n , r , and i are the decimal equivalents of the nominal (or market), real, and inflation rates, respectively. Conversions between real and nominal rates should be made by multiplying and dividing, not by adding and subtracting.